

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
7 September 2001 (07.09.2001)

PCT

(10) International Publication Number  
**WO 01/64674 A1**(51) International Patent Classification<sup>7</sup>: C07D 417/04,  
A61K 31/425, A61P 29/00, C07D 471/04, 417/12, 277/42,  
487/04, 513/04, 417/14KENNIS, Ludo, Edmond, Josephine [BE/BE]; Janssen  
Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse  
(BE).

(21) International Application Number: PCT/EP01/01841

(74) Agent: QUAGHEBEUR, Luc; Janssen Pharmaceutica  
N.V., Turnhoutseweg 30, B-2340 Beerse (BE).

(22) International Filing Date: 20 February 2001 (20.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
00200733.4 1 March 2000 (01.03.2000) EP(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(71) Applicant (*for all designated States except US*):  
JANSSEN PHARMACEUTICA N.V. [BE/BE]; DE  
CORTE, Filip - ext. 3834, Patent Department, Turnhout-  
seweg 30, B-2340 Beerse (BE).(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and





(75) Inventors/Applicants (*for US only*): LOVE, Christo-  
pher [GB/BE]; Janssen Pharmaceutica, Turnhoutseweg 30,  
B-2340 Beerse (BE). VAN WAUWE, Jean, Pierre, Frans  
[BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30,  
B-2340 Beerse (BE). DE BRABANDER, Marc [BE/BE];  
Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340  
Beerse (BE). COOYMANS, Ludwig [BE/BE]; Janssen  
Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse  
(BE). VANDERMAESEN, Nele [BE/BE]; Janssen Phar-  
maceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).

## Published:

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: 2,4-DISUBSTITUTED THIAZOLYL DERIVATIVES



(57) Abstract: This invention concerns the use of a compound of formula (I'), an (N)-oxide, pharmaceutically acceptable addition salt, quaternary amine and stereochemically isomeric form thereof, wherein Q is optionally substituted C<sub>3-6</sub>cycloalkyl, phenyl, naphthyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl; or Q is a radical of formula (b-1, b-2, b-3), wherein X and Y each independently are O, NR<sup>3</sup>, CH<sub>2</sub> or S, with R<sup>3</sup> being hydrogen or C<sub>1-4</sub>alkyl; q is 1 to 4; Z is O or NR<sup>4</sup> with R<sup>4</sup> being hydrogen or C<sub>1-4</sub>alkyl; r is 1 to 3; L is optionally substituted phenyl or L is Het with Het being an optionally substituted five- or six-membered heterocyclic ring or an optionally substituted bicyclic heterocyclic ring; for the manufacture of a medicament for the prevention or the treatment of diseases mediated through cytokines.

2,4-DISUBSTITUTED THIAZOLYL DERIVATIVES

---

5 The present invention is concerned with 2,4-disubstituted thiazolyl derivatives having proinflammatory cytokine production inhibiting properties and adenosine A<sub>3</sub> receptor blocking properties. The invention further relates to methods for their preparation and pharmaceutical compositions comprising them. The invention also relates to the use of 2,4-disubstituted thiazolyl derivatives for the manufacture of a medicament for the prevention or the treatment of warm-blooded animals suffering from diseases mediated through cytokines or diseases mediated through activation of the adenosine A<sub>3</sub> receptor.

- 10 JP 41020220 describes 2-(2-substituted-4-thiazolyl)benzimidazole derivatives as anthelmintics and insecticides.
- J. Prakt. Chem., 1976, 318(5), 875-877 describes the synthesis of pyridyl thiazoles.
- 15 J. Indian. Chem. Soc., 1974, 51(5), 566-568 describes the synthesis and anti-inflammatory activity of some 2-(2-amino-4-thiazolyl)benzothiazoles.
- Fresenius'Z. Anal. Chem., 1977, 288(4), 285 describes the TLC separation of some 2- and 6-[2-amino(and substituted amino)-4-thiazolyl]benzothiazoles.
- 20 Indian J. Chem., 1978, 16B(5), 402-404 describes the synthesis and analgesic, anti-inflammatory activity of 4-(2-amino-4-thiazolyl)isothiazoles.
- WO 97/03073 describes the preparation of thiazolyl triazolothiazoles as anti-ulcer agents and gastric acid secretion inhibitors.
- Indian J. Chem., 1979, 17B(5), 519-521 describes the synthesis of 2-amino-6-benzothiazolyl-2-arylaminothiazoles.
- 25 Indian J. Chem., 1987, 26B(9), 856-860 describes the synthesis and antituberculosis activity of 2-pyrazinyl-2-arylaminothiazoles.
- WO 92/16527 describes the synthesis of 6-methyl-2-pyridyl-2-arylaminothiazoles as agrochemical and horticultural fungicides.
- J. Heterocycl. Chem., 1970, 7(5), 1137-1141 describes the synthesis of pyridyl substituted 2-aminothiazoles.
- 30 DE 3406329 describes the synthesis of 2-pyridinon-2-arylaminothiazole derivatives as inotropic agents.
- J. Chem. Res., Synop., 1998, 12, 742-743, 3329-3347 describes 2-arylmino thiazole derivatives as intermediates to synthesize 5-arylazothiazoles.
- 35 Synth. Commun., 1998, 28(13), 2371-2378 describes the synthesis of 4-(2-furyl)-2-substituted thiazoles utilizing [hydroxy(tosyloxy)iodo]benzene.

Curr. Sci., 1970, 39(18), 417 describes the synthesis of 4-(2'-thienyl) and 4-(2'-furyl)-thiazoles.

DE 4029771 describes the synthesis of N-heteroaryl-2-nitroanilines as pesticides.

- WO 99/32466 describes the preparation of substituted benzenesulfonamide derivatives  
5 as antagonists of the neuropeptide NPY receptor subtype Y5.

Egypt. J. Chem., 1983, 25(2), 187-189 describes the synthesis of sulfamylanilino substituted thiazoles showing bactericidal and fungicidal activity.

Am. Khim. Zh., 1989, 42(10), 657-659 describes the synthesis of  $\delta$ -lactones with heterocyclic substituents.

- 10 Indian J. Chem., Sect. B, 1984, 23B(4), 390-392 describes the synthesis of thiazolylchromones as potential central nervous system agents.

Biol. Zh. Am., 1989, 42(9-10), 956-959 describes the synthesis and activity of unsaturated  $\gamma$ -lactones with thiazole fragments on the growth and development of vegetable crops.


- 15 WO 99/21555 relates to pyridyl substituted thiazolyl compounds having adenosine A<sub>3</sub> receptor antagonistic activity.

WO 99/64418 concerns aryl pyridinyl thiazoles exhibiting inhibition of the human adenosine A<sub>3</sub> receptor activation and of tumor necrosis factor alpha production.

- 20 The compounds of the present invention are distinguishable from the prior art because of their structure, pharmacological activity or potency.

The present invention relates to the use of a compound for the manufacture of a medicament for the prevention or the treatment of diseases mediated through cytokines,

- 25 wherein the compound is a compound of formula




a N-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof,  
wherein

- 30 Q is C<sub>3-6</sub>cycloalkyl, phenyl, naphthyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxy; azido; amino; mono- or di(C<sub>1-6</sub>alkyl)amino; C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>2-6</sub>alkynyl; C<sub>3-6</sub>cycloalkyl; C<sub>1-6</sub>alkyl  
35 substituted with hydroxy, C<sub>1-6</sub>alkyloxy, amino, mono- or di(C<sub>1-4</sub>alkyl)amino;

C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylthio; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl;  
 arylC<sub>1-6</sub>alkyloxy; aryloxy; polyhaloC<sub>1-6</sub>alkyl; polyhalo-C<sub>1-6</sub>alkyloxy; polyhalo-  
 C<sub>1-6</sub>alkylcarbonyl; C<sub>1-4</sub>alkyl-S(=O)<sub>n</sub>- or R<sup>1</sup>HN-S(=O)<sub>n</sub>-;

or

- 5 Q is a radical of formula



wherein X and Y each independently are O, NR<sup>3</sup>, CH<sub>2</sub> or S, with R<sup>3</sup> being  
 hydrogen or C<sub>1-4</sub>alkyl;


q is an integer with value 1 to 4;

- 10 Z is O or NR<sup>4</sup> with R<sup>4</sup> being hydrogen or C<sub>1-4</sub>alkyl;

r is an integer with value 1 to 3;

n is an integer with value 1 or 2;

R<sup>1</sup> represents hydrogen, or a radical of formula



- 15 with A being O, S or a bivalent radical of formula -CR<sup>2a</sup>=N- with CR<sup>2a</sup>  
 attached to N of formula (a-1); and

R<sup>2a</sup> being hydrogen, C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy;

L is phenyl, optionally substituted with up to 4 substituents each independently being  
 selected from halo, hydroxy, amino, cyano, carboxyl, mono-or

- 20 di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or  
 C<sub>1-4</sub>alkyloxy or amino or mono-or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl,  
 C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl,  
 mono-or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-  
 C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-; or

- 25 L is Het;


Het is (i) an optionally substituted five- or six-membered heterocyclic ring containing  
 at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each  
 independently being selected where possible from N, S or O;

- 30 (ii) an optionally substituted five- or six-membered heterocyclic ring containing  
 at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each  
 independently being selected where possible from N, S or O and being fused  
 through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom  
 with another optionally substituted five- or six-membered ring, which

contains, apart from the atoms in common with the first ring, only carbon atoms; the latter ring may be unsaturated, partially unsaturated or saturated;  
 (iii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and at least one heteroatom and being fused through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered heterocyclic ring, which contains, apart from the atoms in common with the first ring, at least one heteroatom; the latter ring may be unsaturated, partially unsaturated or saturated; said bicyclic ring system contains in total from 2 up to 6 heteroatoms, each independently being selected where possible from N, S or O;

wherein Het being a monocyclic ring system may optionally be substituted with up to 4 substituents, and wherein Het being a bicyclic ring system may optionally be substituted with up to 6 substituents, said substituents each independently being selected from halo, hydroxy, amino, cyano, carboxyl, mono- or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono- or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-; aryl is phenyl, optionally substituted with up to five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylthio, cyano, nitro, amino, mono- or di(C<sub>1-6</sub>alkyl)amino.

The present invention also relates to a compound of formula




a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof,  
 wherein

Q is C<sub>3-6</sub>cycloalkyl, phenyl, naphthyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxy; azido; amino; mono- or di(C<sub>1-6</sub>alkyl)amino; C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>2-6</sub>alkynyl; C<sub>3-6</sub>cycloalkyl; C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, amino, mono- or di(C<sub>1-4</sub>alkyl)amino;

C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylthio; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl; arylC<sub>1-6</sub>alkyloxy; aryloxy; polyhaloC<sub>1-6</sub>alkyl; polyhalo-C<sub>1-6</sub>alkyloxy; polyhalo-C<sub>1-6</sub>alkylcarbonyl; C<sub>1-4</sub>alkyl-S(=O)<sub>n</sub>- or R<sup>1</sup>HN-S(=O)<sub>n</sub>-;

or

5 Q is a radical of formula



wherein X and Y each independently are O, NR<sup>3</sup>, CH<sub>2</sub> or S, with R<sup>3</sup> being hydrogen or C<sub>1-4</sub>alkyl;


q is an integer with value 1 to 4;

10 Z is O or NR<sup>4</sup> with R<sup>4</sup> being hydrogen or C<sub>1-4</sub>alkyl;

r is an integer with value 1 to 3;

n is an integer with value 1 or 2;

R<sup>1</sup> represents hydrogen, or a radical of formula



15 with A being O, S or a bivalent radical of formula -CR<sup>2a</sup>=N- with CR<sup>2a</sup> attached to N of formula (a-1); and

R<sup>2a</sup> being hydrogen, C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy;

L is 3-halophenyl, optionally substituted with 1, 2 or 3 substituents each independently being selected from halo, hydroxy, amino, cyano, carboxyl, mono- or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono- or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-; or

25 L is Het;

Het is (i) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each independently being selected where possible from N, S or O;

30 (ii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each independently being selected where possible from N, S or O and being fused through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered ring, which

contains, apart from the atoms in common with the first ring, only carbon atoms; the latter ring may be unsaturated, partially unsaturated or saturated;

(iii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and at least one heteroatom and being fused through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered heterocyclic ring, which contains, apart from the atoms in common with the first ring, at least one heteroatom; the latter ring may be unsaturated, partially unsaturated or saturated; said bicyclic ring system contains in total from 2 up to 6 heteroatoms, each independently being selected where possible from N, S or O;

wherein Het being a monocyclic ring system may optionally be substituted with up to 4 substituents, and wherein Het being a bicyclic ring system may optionally be substituted with up to 6 substituents, said substituents each independently being selected from halo, hydroxy, amino, cyano, carboxyl, mono- or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono- or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-; aryl is phenyl, optionally substituted with up to five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylthio, cyano, nitro, amino, mono- or di(C<sub>1-6</sub>alkyl)amino.

provided that

- 25 - Het is other than optionally substituted isothiazolyl, 2-pyridyl, benzthiazolyl, benzoxazinyl and benzoxazinonyl;
- when Q is phenyl substituted with hydroxy or C<sub>1-6</sub>alkyloxy and carboxy or C<sub>1-6</sub>alkyloxycarbonyl then Het is other than 3-pyridyl or 4-pyridyl;
- when Q is phenyl then Het is other than 2-thienyl, 2-furanyl, 5-bromo-2-benzofuranyl, 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl, 2-benzofuranyl, 5-chloro-2-benzimidazolyl, 2-benzimidazolyl, 3-pyridyl, 4-pyridyl, 6-methyl-thiazolo [3,2-b] [1,2,4]triazol-5-yl, 2,6-dimethyl-thiazolo [3,2-b] [1,2,4]triazol-5-yl or 5,6-dihydro-4,5-dimethyl-2(H)-3-pyranonyl;
- when Q is 2-methyl-phenyl then Het is other than 2-thienyl, 2-benzofuranyl or 3-pyridyl;
- 35 - when Q is 4-methoxy-phenyl then Het is other than 2-furanyl, 2-pyrazinyl, 3-pyridyl, 4-pyridyl, 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl, 1,2-dihydro-6-ethyl-2-oxo-

- 3-cyano-5-pyridyl, 4-(dimethylamino)-1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl, 1,2-dihydro-4-methoxy-6-methyl-2-oxo-3-cyano-5-pyridyl or 3-amino-6-methyl-2(1*H*)-5-pyridinonyl;
- 5 - when Q is 2-methoxy-phenyl then Het is other than 2-pyrazinyl, 5-chloro-2-benzimidazolyl, or 3-pyridyl;
- when Q is 4-chloro-phenyl then Het is other than 2-furanyl, 2-thienyl, 5-chloro-2-benzimidazolyl, 2-pyrazinyl, 3-pyridyl, 4-pyridyl or 5,6-dihydro-4,5-dimethyl-2(H)-3-pyranonyl;
- when Q is 3-chloro-phenyl then Het is other than 2-thienyl, 3-pyridyl or 1,2-dihydro-10 6-methyl-2-oxo-3-cyano-5-pyridyl;
- when Q is 2-chloro-phenyl then Het is other than 2-thienyl;
- when Q is 3-methyl-phenyl then Het is other than 2-thienyl or 3-pyridyl;
- when Q is 2,3-dichloro-phenyl then Het is other than 3-pyridyl;
- when Q is 2-ethoxy-phenyl or 3-methoxy-phenyl then Het is other than 2-pyrazinyl;
- 15 - when Q is 4-bromo-phenyl then Het is other than 2-thienyl, or 5-chloro-2-benzimidazolyl;
- when Q is 4-fluoro-phenyl then Het is other than 4-pyridyl;
- when Q is 1-naphthyl then Het is other than 2-thienyl, or 3-pyridyl;
- when Q is 4-methyl-phenyl then Het is other than 2-furanyl, 2-thienyl, 3-pyridyl, 20 2-pyrazinyl or 5,6-dihydro-4,5-dimethyl-2(H)-3-pyranonyl;
- when Q is 4-ethoxy-phenyl then Het is other than 2-pyrazinyl;
- when Q is 2-naphthyl, 2-carboxy-phenyl, 3-carboxy-phenyl, 4-carboxy-phenyl, 4-amino-phenyl or 3-chloro-2,6-dinitro-4-trifluoromethyl-phenyl then Het is other than 2-thienyl;
- 25 - when Q is 4-benzenesulfonamide then Het is other than 2-furanyl, or 1,2,3,4-tetrahydro-2,4-dioxo-5-pyrimidinyl;
- when Q is N-methyl-4-benzenesulfonamide then Het is other than 3-thienyl;
- when Q is N-butyl-4-benzenesulfonamide then Het is other than 2-furanyl;
- when Q is 2-pyridyl then Het is other than 2-pyrazinyl;
- 30 - when Q is 3-pyridyl then Het is other than 2-thienyl, 3-pyridyl, 4-pyridyl or 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl;
- when Q is 2,4-dichloro-phenyl then Het is other than 2-pyrazinyl, or 4-pyridyl;
- when Q is 4-pyridyl then Het is other than 3-quinolinyl or 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl;
- 35 - when Q is 2,4-dimethoxyphenyl, 3,4-dimethoxyphenyl, 4-hydroxyphenyl, 4-methylthiophenyl, or 4-methylsulfinylphenyl then Het is other than 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl.

The L or Q radical as described above for the compounds of formula (I) or (I') may be attached to the remainder of the molecule of formula (I) or (I') through any ring carbon or heteroatom as appropriate. For example, when Q is pyridyl, it may be 2-pyridyl, 3-pyridyl or 4-pyridyl.

Lines drawn into ring systems indicate that the bond may be attached to any suitable ring atom. When the ring system is a bicyclic ring system, the bond may be attached to any suitable ring atom of either of the two rings.

As used hereinabove or hereinafter C<sub>1-4</sub>alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as methyl, ethyl, propyl, 1-methylethyl, butyl and the like; C<sub>1-6</sub>alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the groups defined for C<sub>1-4</sub>alkyl and pentyl, hexyl, 2-methylbutyl and the like; C<sub>2-6</sub>alkenyl as a group or part of a group defines straight or branched chain hydrocarbon radicals having from 2 to 6 carbon atoms and having 1 double bond such as ethenyl, propenyl, butenyl, pentenyl, hexenyl, 3-methylbutenyl and the like; C<sub>2-6</sub>alkynyl as a group or part of a group defines straight or branched chain hydrocarbon radicals having from 2 to 6 carbon atoms and having 1 triple bond such as ethynyl, propynyl, butynyl, pentynyl, hexynyl, 3-methylbutynyl and the like; C<sub>3-6</sub>cycloalkyl is generic to cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

As used herein before, the term (=O) forms a carbonyl moiety when attached to a carbon atom, a sulfoxide moiety when attached to a sulfur atom and a sulfonyl moiety when two of said terms are attached to a sulfur atom.

The term halo is generic to fluoro, chloro, bromo and iodo. As used in the foregoing and hereinafter, polyhaloC<sub>1-6</sub>alkyl as a group or part of a group is defined as mono- or polyhalosubstituted C<sub>1-6</sub>alkyl, in particular methyl with one or more fluoro atoms, for example, difluoromethyl or trifluoromethyl. In case more than one halogen atoms are attached to an alkyl group within the definition of polyhaloC<sub>1-6</sub>alkyl, they may be the same or different.

When any variable (e.g. R<sup>2a</sup>) occurs more than one time in any constituent, each definition is independent.

It will be appreciated that some of the compounds of formula (I) or (I') and their *N*-oxides, addition salts, quaternary amines and stereochemically isomeric forms may contain one or more centers of chirality and exist as stereochemically isomeric forms.

5

The term "stereochemically isomeric forms" as used hereinbefore or hereinafter defines all the possible stereoisomeric forms which the compounds of formula (I) or (I') and their *N*-oxides, addition salts, quaternary amines or physiologically functional derivatives may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure as well as each of the individual isomeric forms of formula (I) or (I') and their *N*-oxides, salts, solvates, quaternary amines substantially free, *i.e.* associated with less than 10%, preferably less than 5%, in particular less than 2% and most preferably less than 1% of the other isomers. Stereochemically isomeric forms of the compounds of formula (I) or (I') are obviously intended to be embraced within the scope of this invention.

For therapeutic use, salts of the compounds of formula (I) or (I') are those wherein the counterion is pharmaceutically acceptable. However, salts of acids and bases which are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

25

The pharmaceutically acceptable acid and base addition salts as mentioned hereinabove or hereinafter are meant to comprise the therapeutically active non-toxic acid and base addition salt forms which the compounds of formula (I) or (I') are able to form. The pharmaceutically acceptable acid addition salts can conveniently be obtained by treating the base form with such appropriate acid. Appropriate acids comprise, for example, inorganic acids such as hydrohalic acids, *e.g.* hydrochloric or hydrobromic acid, sulfuric, nitric, phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic (*i.e.* ethanedioic), malonic, succinic (*i.e.* butanedioic acid), maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

35

Conversely said salt forms can be converted by treatment with an appropriate base into the free base form.

The compounds of formula (I) or (I') containing an acidic proton may also be  
5 converted into their non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. primary, secondary and tertiary aliphatic and aromatic amines such as methylamine,  
10 ethylamine, propylamine, isopropylamine, the four butylamine isomers, dimethylamine, diethylamine, diethanolamine, dipropylamine, diisopropylamine, di-n-butylamine, pyrrolidine, piperidine, morpholine, trimethylamine, triethylamine, tripropylamine, quinuclidine, pyridine, quinoline and isoquinoline; the benzathine, *N*-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for  
15 example, arginine, lysine and the like.











Conversely the salt form can be converted by treatment with acid into the free acid form.

The term addition salt as used hereinabove also comprises the solvates which the  
20 compounds of formula (I) or (I') as well as the salts thereof, are able to form. Such solvates are for example hydrates, alcoholates and the like.

The term "quaternary amine" as used hereinbefore defines the quaternary ammonium salts which the compounds of formula (I) or (I') are able to form by reaction between a  
25 basic nitrogen of a compound of formula (I) or (I') and an appropriate quaternizing agent, such as, for example, an optionally substituted alkylhalide, arylhalide or arylalkylhalide, e.g. methyl iodide or benzyl iodide. Other reactants with good leaving groups may also be used, such as alkyl trifluoromethanesulfonates, alkyl methanesulfonates, and alkyl *p*-toluenesulfonates. A quaternary amine has a positively  
30 charged nitrogen. Pharmaceutically acceptable counterions include for example chloro, bromo, iodo, trifluoroacetate and acetate. The counterion of choice can be made using ion exchange resin columns.

Some of the compounds of formula (I) or (I') may also exist in their tautomeric form.  
35 Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.

In particular, the radical Het as defined hereinabove may be a radical of formula



with A<sub>1</sub>, B<sub>1</sub>, C<sub>1</sub>, D<sub>1</sub> and E<sub>1</sub>, each independently being selected where possible from CH, N, NH, O or S, provided that from 1 up to 4 heteroatoms are present, and wherein each C or N atom, where possible, may optionally be substituted with halo, hydroxy, amino, cyano, carboxyl, mono-or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono-or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono-or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-, said substituents being limited to a total of 4, and wherein each dotted line may represent, where possible, an additional bond, provided that two double bonds are present;

with A<sub>2</sub>, B<sub>2</sub>, C<sub>2</sub>, D<sub>2</sub>, E<sub>2</sub> and F<sub>2</sub>, each independently being selected where possible from CH, N, O or S, provided that from 1 up to 4 heteroatoms are present, and wherein each C or N atom, where possible, may optionally be substituted with halo, hydroxy, amino, cyano, carboxyl, mono-or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono-or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono-or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-, said substituents being limited to a total of 4, and wherein each dotted line may represent, where possible, an additional bond, provided that at least two double bonds are present;

with A<sub>3</sub> and E<sub>3</sub>, each independently being selected where possible from C, CH or N, and B<sub>3</sub>, C<sub>3</sub> and D<sub>3</sub>, each independently and where possible being selected from CH, CH<sub>2</sub>, N, NH, O or S, and F<sub>3</sub>, G<sub>3</sub>, H<sub>3</sub> and I<sub>3</sub>, each independently and where possible being selected from CH<sub>2</sub> or CH, provided that from 1 up to 4 heteroatoms are present, and wherein each C or N atom, where possible, may optionally be substituted with halo, hydroxy, amino, cyano, carboxyl, mono-or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono-or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono-or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-, said substituents being limited to a total of 6, and wherein each dotted line may represent, where possible, an additional bond, provided that the five-membered ring contains two double bonds;

with A<sub>4</sub> and E<sub>4</sub>, each independently being selected where possible from C, CH or N, and B<sub>4</sub>, C<sub>4</sub> and D<sub>4</sub>, each independently and where possible being selected from CH, CH<sub>2</sub>, N, NH, O or S, and F<sub>4</sub>, G<sub>4</sub>, and H<sub>4</sub>, each independently and where possible being selected from CH<sub>2</sub> or CH, provided that from 1 up to 4 heteroatoms are present, and wherein each C or N atom, where possible, may optionally be substituted with halo, hydroxy, amino, cyano, carboxyl, mono-or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono-or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono-or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-, said substituents being limited to a total of 6, and wherein each dotted line may represent, where possible, an additional bond, provided that the five-membered ring consisting of A<sub>4</sub>-B<sub>4</sub>-C<sub>4</sub>-D<sub>4</sub>-E<sub>4</sub> contains two double bonds;

with A<sub>5</sub> and F<sub>5</sub>, each independently being selected where possible from C, CH or N, and B<sub>5</sub>, C<sub>5</sub>, D<sub>5</sub> and E<sub>5</sub>, each independently and where possible being selected from CH, CH<sub>2</sub>, N, O or S, and G<sub>5</sub>, H<sub>5</sub>, I<sub>5</sub> and J<sub>5</sub>, each independently and where possible being selected from CH<sub>2</sub> or CH, provided that from 1 up to 4 heteroatoms are present, and wherein each C or N atom, where possible, may optionally be substituted with halo, hydroxy, amino, cyano, carboxyl, mono-or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono-or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono-or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-, said substituents being limited to a total of 6, and

wherein each dotted line may represent, where possible, an additional bond, provided that the six-membered ring consisting of A<sub>5</sub>-B<sub>5</sub>-C<sub>5</sub>-D<sub>5</sub>-E<sub>5</sub>-F<sub>5</sub> contains at least two double bonds;

- with A<sub>6</sub> and F<sub>6</sub>, each independently being selected where possible from C, CH or N, and B<sub>6</sub>, C<sub>6</sub>, D<sub>6</sub> and E<sub>6</sub>, each independently and where possible being selected from CH, CH<sub>2</sub>, N, O or S, and G<sub>6</sub>, H<sub>6</sub> and I<sub>6</sub>, each independently and where possible being selected from CH<sub>2</sub> or CH, provided that from 1 up to 4 heteroatoms are present, and wherein each C or N atom, where possible, may optionally be substituted with halo, hydroxy, amino, cyano, carboxyl, mono-or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono-or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono-or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-, said substituents being limited to a total of 6, and wherein each dotted line may represent, where possible, an additional bond, provided that the six-membered ring contains at least two double bonds;

- with A<sub>7</sub> and E<sub>7</sub>, each independently being selected where possible from C, CH or N, and B<sub>7</sub>, C<sub>7</sub> and D<sub>7</sub>, each independently and where possible being selected from CH, CH<sub>2</sub>, N, NH, O or S, and F<sub>7</sub>, G<sub>7</sub>, H<sub>7</sub> and I<sub>7</sub>, each independently and where possible being selected from CH, CH<sub>2</sub>, N, NH, O or S, provided that the bicyclic ring contains in total from 2 up to 6 heteroatoms with at least one heteroatom in the five-membered ring and at least one heteroatom in the remainder, i.e. F<sub>7</sub>-G<sub>7</sub>-H<sub>7</sub>-I<sub>7</sub>, of the fused six-membered ring, and wherein each C or N atom, where possible, may optionally be substituted with halo, hydroxy, amino, cyano, carboxyl, mono-or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono-or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono-or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-, said substituents being limited to a total of 6, and wherein each dotted line may represent, where possible, an additional bond, provided that the five-membered ring contains two double bonds;

- with A<sub>8</sub> and E<sub>8</sub>, each independently being selected where possible from C, CH or N, and B<sub>8</sub>, C<sub>8</sub>, and D<sub>8</sub>, each independently and where possible being selected from CH, CH<sub>2</sub>, N, NH, O or S, and F<sub>8</sub>, G<sub>8</sub>, and H<sub>8</sub>, each independently and where possible being selected from CH, CH<sub>2</sub>, N, NH, O or S, provided that the bicyclic ring contains in total from 2 up to 6 heteroatoms with at least one heteroatom in the five-membered ring consisting of A<sub>8</sub>-B<sub>8</sub>-C<sub>8</sub>-D<sub>8</sub>-E<sub>8</sub> and at least one heteroatom in the remainder, i.e. F<sub>8</sub>-G<sub>8</sub>-


- H<sub>8</sub>, of the other, fused five-membered ring, and wherein each C or N atom, where possible, may optionally be substituted with halo, hydroxy, amino, cyano, carboxyl, mono- or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono- or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-, said substituents being limited to a total of 6, and wherein each dotted line may represent, where possible, an additional bond, provided that the five-membered ring consisting of A<sub>8</sub>-B<sub>8</sub>-C<sub>8</sub>-D<sub>8</sub>-E<sub>8</sub> contains two double bonds;
- with A<sub>9</sub> and F<sub>9</sub>, each independently being selected where possible from C, CH or N, and B<sub>9</sub>, C<sub>9</sub>, D<sub>9</sub> and E<sub>9</sub>, each independently and where possible being selected from CH, CH<sub>2</sub>, N, O or S, and G<sub>9</sub>, H<sub>9</sub>, I<sub>9</sub> and J<sub>9</sub>, each independently and where possible being selected from CH, CH<sub>2</sub>, N, NH, O or S, provided that the bicyclic ring contains in total from 2 up to 6 heteroatoms with at least one heteroatom in the six-membered ring consisting of A<sub>9</sub>-B<sub>9</sub>-C<sub>9</sub>-D<sub>9</sub>-E<sub>9</sub>-F<sub>9</sub> and at least one heteroatom in the remainder, i.e. G<sub>9</sub>-H<sub>9</sub>-I<sub>9</sub>-J<sub>9</sub>, of the other, fused six-membered ring, and wherein each C or N atom, where possible, may optionally be substituted with halo, hydroxy, amino, cyano, carboxyl, mono- or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono- or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-, said substituents being limited to a total of 6, and wherein each dotted line may represent, where possible, an additional bond, provided that the six-membered ring consisting of A<sub>9</sub>-B<sub>9</sub>-C<sub>9</sub>-D<sub>9</sub>-E<sub>9</sub>-F<sub>9</sub> contains at least two double bonds;
- with A<sub>10</sub> and F<sub>10</sub>, each independently being selected where possible from C, CH or N, and B<sub>10</sub>, C<sub>10</sub>, D<sub>10</sub> and E<sub>10</sub>, each independently and where possible being selected from CH, CH<sub>2</sub>, N, O or S, and G<sub>10</sub>, H<sub>10</sub> and I<sub>10</sub>, each independently and where possible being selected from CH, CH<sub>2</sub>, N, NH, O or S, provided that the bicyclic ring contains in total from 2 up to 6 heteroatoms with at least one heteroatom in the six-membered ring and at least one heteroatom in the remainder, i.e. G<sub>10</sub>-H<sub>10</sub>-I<sub>10</sub>, of the fused five-membered ring, and wherein each C or N atom, where possible, may optionally be substituted with halo, hydroxy, amino, cyano, carboxyl, mono- or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono- or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono- or

di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-, said substituents being limited to a total of 6, and wherein each dotted line may represent, where possible, an additional bond, provided that the six-membered ring contains at least two double bonds.

More in particular, the radical Het as defined hereinabove may be a monocyclic heterocycle comprising furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, isoxazolyl, isothiazolyl, pyrazolyl, oxadiazolyl, thiadiazolyl, triazolyl, 1-pyridyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, pyranyl, triazinyl, tetrazolyl, with each monocyclic heterocycle optionally substituted with, where possible, one, two, three or four substituents selected from halo, hydroxy, amino, cyano, carboxyl, mono- or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with C<sub>1-4</sub>alkyloxy or amino or mono- or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxy carbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-; or Het may also represent a bicyclic heterocycle comprising benzofuranyl, benzothienyl, benzthiazolyl, benzoxazinyl, benzoxazinonyl, indoliziny, indolyl, isoindolyl, benzoxazolyl, benzimidazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, benzopyrazolyl, benzoxadiazolyl, benzothiadiazolyl, benzotriazolyl, naphthalenyl, quinoliny, isoquinoliny, cinnoliny, phthalazinyl, quinoxaliny, quinazoliny, naphthiridiny, benzopyranyl, pyrrolopyridyl, thienopyridyl, furopyridyl, isothiazolopyridyl, thiazolopyridyl, isoxazolopyridyl, oxazolopyridyl, pyrazolopyridyl, imidazopyridyl, pyrrolopyrazinyl, thienopyrazinyl, furopyrazinyl, isothiazolopyrazinyl, thiazolopyrazinyl, isoxazolopyrazinyl, oxazolopyrazinyl, pyrazolopyrazinyl, imidazopyrazinyl, pyrrolopyrimidinyl, thienopyrimidinyl, furopyrimidinyl, isothiazolopyrimidinyl, thiazolopyrimidinyl, isoxazolopyrimidinyl, oxazolopyrimidinyl, pyrazolopyrimidinyl, imidazopyrimidinyl, pyrrolopyridazinyl, thienopyridazinyl, furopyridazinyl, isothiazolopyridazinyl, thiazolopyridazinyl, isoxazolopyridazinyl, oxazolopyridazinyl, pyrazolopyridazinyl, imidazopyridazinyl, oxadiazolopyridyl, thiadiazolopyridyl, triazolopyridyl, oxadiazolopyrazinyl, thiadiazolopyrazinyl, triazolopyrazinyl, oxadiazolopyrimidinyl, thiadiazolopyrimidinyl, triazolopyrimidinyl, oxadiazolopyridazinyl, thiadiazolopyridazinyl, triazolopyridazinyl, imidazooxazolyl, imidazothiazolyl, imidazoimidazolyl, isoxazolotriazinyl, isothiazolotriazinyl, pyrazolotriazinyl, oxazolotriazinyl, thiazolotriazinyl, imidazotriazinyl, oxadiazolotriazinyl, thiadiazolotriazinyl, triazolotriazinyl, with each bicyclic heterocycle optionally substituted with, where possible, up to 6 substituents selected from

- halo, hydroxy, amino, cyano, carboxyl, mono-or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with C<sub>1-4</sub>alkyloxy or amino or mono-or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono-or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-,
- 5 C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-.

- In particular, the present invention relates to the use of a compound for the manufacture of a medicament for the prevention or the treatment of diseases mediated through
- 10 cytokines, wherein the compound is a compound of formula




a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof, wherein

- 15 Q is C<sub>3-6</sub>cycloalkyl, phenyl, naphthyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxy; azido; amino; mono-
- 20 or di(C<sub>1-6</sub>alkyl)amino; C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>2-6</sub>alkynyl; C<sub>3-6</sub>cycloalkyl; C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, amino, mono-or di(C<sub>1-4</sub>alkyl)amino; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylthio; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl; arylC<sub>1-6</sub>alkyloxy; aryloxy; polyhaloC<sub>1-6</sub>alkyl; polyhalo-C<sub>1-6</sub>alkyloxy; polyhalo-C<sub>1-6</sub>alkylcarbonyl; C<sub>1-4</sub>alkyl-S(=O)<sub>n</sub>- or R<sup>1</sup>HN-S(=O)<sub>n</sub>-;

or

- 25 Q is a radical of formula



wherein X and Y each independently are O, NR<sup>3</sup>, CH<sub>2</sub> or S, with R<sup>3</sup> being hydrogen or C<sub>1-4</sub>alkyl;

q is an integer with value 1 to 4;


- 30 Z is O or NR<sup>4</sup> with R<sup>4</sup> being hydrogen or C<sub>1-4</sub>alkyl;

r is an integer with value 1 to 3;

n is an integer with value 1 or 2;

R<sup>1</sup> represents hydrogen, or a radical of formula

-17-



with A being O, S or a bivalent radical of formula  $-\text{CR}^{2a}=\text{N}-$  with  $\text{CR}^{2a}$  attached to N of formula (a-1); and

$\text{R}^{2a}$  being hydrogen,  $\text{C}_{1-6}$ alkyl or  $\text{C}_{1-6}$ alkyloxy;

5 L is Het;

Het is (i) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each independently being selected where possible from N, S or O;

10 (ii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each independently being selected where possible from N, S or O and being fused through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered ring, which contains, apart from the atoms in common with the first ring, only carbon

15 atoms; the latter ring may be unsaturated, partially unsaturated or saturated;

(iii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and at least one heteroatom and being fused through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered heterocyclic

20 ring, which contains, apart from the atoms in common with the first ring, at least one heteroatom; the latter ring may be unsaturated, partially unsaturated or saturated; said bicyclic ring system contains in total from 2 up to 6 heteroatoms, each independently being selected where possible from N, S or O;


25 wherein Het being a monocyclic ring system may optionally be substituted with up to 4 substituents, and wherein Het being a bicyclic ring system may optionally be substituted with up to 6 substituents, said substituents each independently being selected from halo, hydroxy, amino, cyano, carboxyl, mono- or di( $\text{C}_{1-4}$ alkyl)amino,  $\text{C}_{1-6}$ alkyl,  $\text{C}_{1-6}$ alkyl substituted with hydroxy or

30  $\text{C}_{1-4}$ alkyloxy or amino or mono- or di( $\text{C}_{1-4}$ alkyl)amino, polyhalo $\text{C}_{1-6}$ alkyl,  $\text{C}_{1-6}$ alkyloxy,  $\text{C}_{1-6}$ alkyloxycarbonyl,  $\text{C}_{1-6}$ alkylcarbonyloxy, aminocarbonyl, mono- or di( $\text{C}_{1-6}$ alkyl)aminocarbonyl,  $\text{C}_{1-6}$ alkyl- $\text{C}(=\text{O})-\text{NH}-$ ,  $\text{C}_{1-6}$ alkyloxy- $\text{C}(=\text{O})-\text{NH}-$ ,  $\text{H}_2\text{N}-\text{C}(=\text{O})-\text{NH}-$  or mono- or di( $\text{C}_{1-4}$ alkyl)amino- $\text{C}(=\text{O})-\text{NH}-$ ;

-18-

aryl is phenyl, optionally substituted with up to five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylthio, cyano, nitro, amino, mono-or di(C<sub>1-6</sub>alkyl)amino; provided that Het is other than optionally substituted isothiazolyl, 2-pyridyl, benzthiazolyl, benzoxazinyl and benzoxazinonyl.

More in particular, the present invention relates to the use of a compound for the manufacture of a medicament for the prevention or the treatment of diseases mediated through cytokines, wherein the compound is a compound of formula




a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof, wherein

Q is C<sub>3-6</sub>cycloalkyl, phenyl, naphthyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxy; azido; amino; mono-or di(C<sub>1-6</sub>alkyl)amino; C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>2-6</sub>alkynyl; C<sub>3-6</sub>cycloalkyl; C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, amino, mono-or di(C<sub>1-4</sub>alkyl)amino; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylthio; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxy carbonyl; arylC<sub>1-6</sub>alkyloxy; aryloxy; polyhaloC<sub>1-6</sub>alkyl; polyhalo-C<sub>1-6</sub>alkyloxy; polyhalo-C<sub>1-6</sub>alkylcarbonyl; C<sub>1-4</sub>alkyl-S(=O)<sub>n</sub>- or R<sup>1</sup>HN-S(=O)<sub>n</sub>-;

or

Q is a radical of formula



wherein X and Y each independently are O, NR<sup>3</sup>, CH<sub>2</sub> or S, with R<sup>3</sup> being hydrogen or C<sub>1-4</sub>alkyl;


q is an integer with value 1 to 4;

Z is O or NR<sup>4</sup> with R<sup>4</sup> being hydrogen or C<sub>1-4</sub>alkyl;

r is an integer with value 1 to 3;

n is an integer with value 1 or 2;

R<sup>1</sup> represents hydrogen, or a radical of formula



with A being O, S or a bivalent radical of formula  $-\text{CR}^{2a}=\text{N}-$  with  $\text{CR}^{2a}$  attached to N of formula (a-1); and

$\text{R}^{2a}$  being hydrogen,  $\text{C}_{1-6}$ alkyl or  $\text{C}_{1-6}$ alkyloxy;

5 L is Het;

Het is (i) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each independently being selected where possible from N, S or O;

10 (ii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each independently being selected where possible from N, S or O and being fused through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered ring, which contains, apart from the atoms in common with the first ring, only carbon atoms; the latter ring may be unsaturated, partially unsaturated or saturated;

15 (iii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and at least one heteroatom and being fused through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered heterocyclic ring, which contains, apart from the atoms in common with the first ring, at least one heteroatom; the latter ring may be unsaturated, partially unsaturated or saturated; said bicyclic ring system contains in total from 2 up to 6 heteroatoms, each independently being selected where possible from N, S or O;

25 wherein Het being a monocyclic ring system may optionally be substituted with up to 4 substituents, and wherein Het being a bicyclic ring system may optionally be substituted with up to 6 substituents, said substituents each independently being selected from halo, hydroxy, amino, cyano, carboxyl, mono- or di( $\text{C}_{1-4}$ alkyl)amino,  $\text{C}_{1-6}$ alkyl,  $\text{C}_{1-6}$ alkyl substituted with hydroxy or


30  $\text{C}_{1-4}$ alkyloxy or amino or mono- or di( $\text{C}_{1-4}$ alkyl)amino, polyhalo $\text{C}_{1-6}$ alkyl,  $\text{C}_{1-6}$ alkyloxy,  $\text{C}_{1-6}$ alkyloxycarbonyl,  $\text{C}_{1-6}$ alkylcarbonyloxy, aminocarbonyl, mono- or di( $\text{C}_{1-6}$ alkyl)aminocarbonyl,  $\text{C}_{1-6}$ alkyl- $\text{C}(=\text{O})-\text{NH}-$ ,  $\text{C}_{1-6}$ alkyloxy- $\text{C}(=\text{O})-\text{NH}-$ ,  $\text{H}_2\text{N}-\text{C}(=\text{O})-\text{NH}-$  or mono- or di( $\text{C}_{1-4}$ alkyl)amino- $\text{C}(=\text{O})-\text{NH}-$ ;

aryl is phenyl, optionally substituted with up to five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylthio, cyano, nitro, amino, mono-or di(C<sub>1-6</sub>alkyl)amino; provided that

- 5 - Het is other than optionally substituted isothiazolyl, 2-pyridyl, benzthiazolyl, benzoxazinyl and benzoxazinonyl.
- when Q is phenyl substituted with hydroxy or C<sub>1-6</sub>alkyloxy and carboxy or C<sub>1-6</sub>alkyloxycarbonyl then Het is other than 3-pyridyl or 4-pyridyl;
- when Q is phenyl then Het is other than 2-thienyl, 2-furanyl, 5-bromo-2-benzofuranyl, 10 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl, 2-benzofuranyl, 5-chloro-2-benzimidazolyl, 2-benzimidazolyl, 3-pyridyl, 4-pyridyl, 6-methyl-thiazolo [3,2-b] [1,2,4]triazol-5-yl, 2,6-dimethyl-thiazolo [3,2-b] [1,2,4]triazol-5-yl or 5,6-dihydro-4,5-dimethyl-2(H)-3-pyranonyl;
- when Q is 2-methyl-phenyl then Het is other than 2-thienyl, 2-benzofuranyl or 15 3-pyridyl;
- when Q is 4-methoxy-phenyl then Het is other than 2-furanyl, 2-pyrazinyl, 3-pyridyl, 4-pyridyl, 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl, 1,2-dihydro-6-ethyl-2-oxo-3-cyano-5-pyridyl, 4-(dimethylamino)-1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl, 1,2-dihydro-4-methoxy-6-methyl-2-oxo-3-cyano-5-pyridyl or 3-amino-6-methyl-2(1H)-5-pyridinonyl; 20
- when Q is 2-methoxy-phenyl then Het is other than 2-pyrazinyl, 5-chloro-2-benzimidazolyl, or 3-pyridyl;
- when Q is 4-chloro-phenyl then Het is other than 2-furanyl, 2-thienyl, 5-chloro-2-benzimidazolyl, 2-pyrazinyl, 3-pyridyl, 4-pyridyl or 5,6-dihydro-4,5-dimethyl-2(H)-3-pyranonyl; 25
- when Q is 3-chloro-phenyl then Het is other than 2-thienyl, 3-pyridyl or 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl;
- when Q is 2-chloro-phenyl then Het is other than 2-thienyl;
- when Q is 3-methyl-phenyl then Het is other than 2-thienyl or 3-pyridyl;
- 30 - when Q is 2,3-dichloro-phenyl then Het is other than 3-pyridyl;
- when Q is 2-ethoxy-phenyl or 3-methoxy-phenyl then Het is other than 2-pyrazinyl;
- when Q is 4-bromo-phenyl then Het is other than 2-thienyl, or 5-chloro-2-benzimidazolyl;
- when Q is 4-fluoro-phenyl then Het is other than 4-pyridyl;
- 35 - when Q is 1-naphthyl then Het is other than 2-thienyl, or 3-pyridyl;
- when Q is 4-methyl-phenyl then Het is other than 2-furanyl, 2-thienyl, 3-pyridyl, 2-pyrazinyl or 5,6-dihydro-4,5-dimethyl-2(H)-3-pyranonyl;

- when Q is 4-ethoxy-phenyl then Het is other than 2-pyrazinyl;
- when Q is 2-naphthyl, 2-carboxy-phenyl, 3-carboxy-phenyl, 4-carboxy-phenyl, 4-amino-phenyl or 3-chloro-2,6-dinitro-4-trifluoromethyl-phenyl then Het is other than 2-thienyl;
- 5 - when Q is 4-benzenesulfonamide then Het is other than 2-furanyl, or 1,2,3,4-tetrahydro-2,4-dioxo-5-pyrimidinyl;
- when Q is N-methyl-4-benzenesulfonamide then Het is other than 3-thienyl;
- when Q is N-butyl-4-benzenesulfonamide then Het is other than 2-furanyl;
- when Q is 2-pyridyl then Het is other than 2-pyrazinyl;
- 10 - when Q is 3-pyridyl then Het is other than 2-thienyl, 3-pyridyl, 4-pyridyl or 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl;
- when Q is 2,4-dichloro-phenyl then Het is other than 2-pyrazinyl, or 4-pyridyl;
- when Q is 4-pyridyl then Het is other than 3-quinoliny or 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl;
- 15 - when Q is 2,4-dimethoxyphenyl, 3,4-dimethoxyphenyl, 4-hydroxyphenyl, 4-methylthiophenyl, or 4-methylsulfinylphenyl then Het is other than 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl.

The present invention also relates to the use of a compound for the manufacture of a medicament for the prevention or the treatment of diseases mediated through cytokines, wherein the compound is a compound of formula



a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof,


wherein

Q is phenyl, naphthyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxy; azido; amino; mono- or di(C<sub>1-6</sub>alkyl)amino; C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>2-6</sub>alkynyl; C<sub>3-6</sub>cycloalkyl; C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, amino, mono- or di(C<sub>1-4</sub>alkyl)amino; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylthio; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxy carbonyl; arylC<sub>1-6</sub>alkyloxy; aryloxy; polyhaloC<sub>1-6</sub>alkyl; polyhalo-C<sub>1-6</sub>alkyloxy; polyhalo-C<sub>1-6</sub>alkylcarbonyl; C<sub>1-4</sub>alkyl-S(=O)<sub>n</sub>- or R<sup>1</sup>HN-S(=O)<sub>n</sub>-;

or

Q is a radical of formula

-22-



wherein X and Y each independently are O,  $\text{NR}^3$ ,  $\text{CH}_2$  or S, with  $\text{R}^3$  being hydrogen or  $\text{C}_{1-4}$ alkyl;


q is an integer with value 1 to 4;

5 Z is O or  $\text{NR}^4$  with  $\text{R}^4$  being hydrogen or  $\text{C}_{1-4}$ alkyl;

r is an integer with value 1 to 3;

n is an integer with value 1 or 2;

$\text{R}^1$  represents hydrogen, or a radical of formula



10 with A being O, S or a bivalent radical of formula  $-\text{CR}^{2a}=\text{N}-$  with  $\text{CR}^{2a}$  attached to N of formula (a-1); and

$\text{R}^{2a}$  being hydrogen,  $\text{C}_{1-6}$ alkyl or  $\text{C}_{1-6}$ alkyloxy;

L is phenyl, optionally substituted with up to 4 substituents each independently being selected from halo, hydroxy, amino, mono or di( $\text{C}_{1-4}$ alkyl)amino,  $\text{C}_{1-6}$ alkyl,

15 polyhalo $\text{C}_{1-6}$ alkyl or  $\text{C}_{1-6}$ alkyloxy; or

L is Het;

Het is (i) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each independently being selected where possible from N, S or O;

20 (ii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each independently being selected where possible from N, S or O and being fused through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered ring, which contains, apart from the atoms in common with the first ring, only carbon atoms; the latter ring may be unsaturated, partially unsaturated or saturated;

25 (iii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and at least one heteroatom and being fused through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered heterocyclic ring, which contains, apart from the atoms in common with the first ring, at least one heteroatom; the latter ring may be unsaturated, partially unsaturated or saturated; said bicyclic ring system contains in total from 2

30

up to 6 heteroatoms, each independently being selected where possible from N, S or O;

- wherein Het being a monocyclic ring system may optionally be substituted with up to 4 substituents, and wherein Het being a bicyclic ring system may optionally be substituted with up to 6 substituents, said substituents each independently being selected from halo, hydroxy, amino, mono or di(C<sub>1-6</sub>alkyl)amino, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy; aryl is phenyl, optionally substituted with up to five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylthio, cyano, nitro, amino, mono-or di(C<sub>1-6</sub>alkyl)amino.

- An interesting group comprises those compounds of formula (I) or (I') wherein L is Het and Het is defined as hereinabove provided that Het is other than benzimidazolyl; benzofuranyl; thiazolotriazolyl; quinoliny; pyrazinyl; dioxypyrimidinyl; pyrimidinyl; pyridazinyl; pyranonyl; thienyl; furanyl; a 5 or 6-membered heterocyclic group containing one nitrogen atom such as for example pyridyl.

- Also an interesting group comprises those compounds of formula (I) or (I') wherein L is Het and Het being a monocyclic ring system may optionally be substituted with up to 4 substituents, or Het being a bicyclic ring system may optionally be substituted with up to 6 substituents, said substituents each independently being selected from halo, hydroxy, amino, mono or di(C<sub>1-6</sub>alkyl)amino, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy.

- A further interesting group comprises those compounds of formula (I) or (I') wherein L is imidazolyl, imidazothiazolyl, pyrimidinyl, thienyl, thiazolyl, furanyl, 3-pyridyl, 4-pyridyl, pyrazolyl, indolyl, indazolyl, quinoliny, benzofuranyl, pyrrolopyridyl, imidazopyridyl, imidazopyrazinyl, imidazopyrimidinyl, imidazopyridazinyl, pyrazolopyridyl, with each heterocycle optionally substituted with one, two, three or four substituents selected from halo, amino, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl, aminocarbonyl or C<sub>1-6</sub>alkyl-C(=O)-NH-.

- Still another interesting group includes those compounds of formula (I) or (I') wherein L is 3-pyridyl, 4-pyridyl, thiazolyl, pyrazolyl, indolyl, indazolyl, quinoliny, benzofuranyl, pyrrolopyridyl, imidazopyridyl, imidazopyrazinyl, imidazopyrimidinyl, imidazopyridazinyl, pyrazolopyridyl, with each heterocycle optionally substituted with one, two, three or four substituents selected from halo, amino, or C<sub>1-6</sub>alkyl.

Yet a further interesting group comprises those compounds of formula (I) or (I') wherein L is imidazolyl, imidazothiazolyl, pyrimidinyl, pyrazolyl, indolyl, indazolyl, pyrrolopyridyl, imidazopyridyl, imidazopyrazinyl, imidazopyrimidinyl, imidazopyridazinyl, pyrazolopyridyl, with each heterocycle optionally substituted with one, two, three or four substituents selected from halo, amino, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl, aminocarbonyl or C<sub>1-6</sub>alkyl-C(=O)-NH-.

Again an interesting group comprises those compounds of formula (I) or (I') wherein L is Het and Het is as defined hereinabove provided that Het is other than pyrazolyl, benzofuranyl, 2-imidazo[1,2-a]pyridyl, imidazopyridazinyl, indazolyl, pyrazinyl, 4-pyrimidinyl, thiazolyl, imidazolyl.

Also an interesting group comprises those compounds of formula (I) or (I') wherein L is Het and Het is as defined hereinabove provided that Het is other than pyrazolyl, benzofuranyl, 2-imidazo[1,2-a]pyridyl, imidazopyridazinyl, indazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, thiazolyl, imidazolyl, benzimidazolyl, thiazolotriazolyl, quinoliny, dioxypyrimidinyl, pyranonyl, a 5 or 6-membered heterocyclic group containing one nitrogen atom, thienyl, furanyl.

Again an interesting group comprises those compounds of formula (I) or (I') wherein L is Het and Het is indolyl, 3-imidazo[1,2-a]pyridyl, 3-imidazo[1,5-a]pyridyl, 3-pyridyl, quinoliny, imidazopyrimidinyl, imidazopyrazinyl, imidazothiazolyl, 5-pyrimidinyl, furanyl, thiazolyl, imidazolyl, pyrrolopyridyl, pyrazolopyridyl.

A further interesting group comprises those compounds of formula (I) or (I') wherein L is Het and Het is indolyl, 3-imidazo[1,2-a]pyridyl, 3-imidazo[1,5-a]pyridyl, imidazopyrimidinyl, imidazopyrazinyl, imidazothiazolyl, pyrrolopyridyl, pyrazolopyridyl.


Further preferred compounds are those compounds of formula (I) or (I') wherein L is Het and Het is 3-imidazo[1,2-a]pyridyl, 3-imidazo[1,5-a]pyridyl, imidazothiazolyl, 5-pyrimidinyl, substituted 3- or 4-pyridyl.

Yet further preferred compounds are those compounds of formula (I) or (I') wherein L is 3-imidazo[1,2-a]pyridyl, 3-imidazo[1,5-a]pyridyl, imidazothiazolyl, 3-pyridyl or pyrrolopyridyl.

Also preferred compounds are those compounds of formula (I) or (I') wherein L is 3-fluorophenyl or 3,5-difluorophenyl.

- 5 Also preferred are those compounds of formula (I) or (I') wherein L is Het and Het is as described hereinabove provided that the atom(s) adjacent to the atom with which Het is linked to the remainder of the molecule of formula (I) and which does (do) not form part of both rings in case of a bicyclic heterocycle, is (are) other than nitrogen.
- 10 Again preferred compounds are those compounds of formula (I) or (I') wherein L is 3-halophenyl.

- Also an interesting group comprises those compounds of formula (I) or (I') wherein Q is phenyl, naphthyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl or imidazopyridyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; azido; amino; mono- or di(C<sub>1-6</sub>alkyl)amino; C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>2-6</sub>alkynyl; C<sub>3-6</sub>cycloalkyl; C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, amino, mono- or di(C<sub>1-4</sub>alkyl)amino; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylthio; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl; arylC<sub>1-6</sub>alkyloxy; aryloxy; polyhaloC<sub>1-6</sub>alkyl; polyhaloC<sub>1-6</sub>alkyloxy; polyhaloC<sub>1-6</sub>alkylcarbonyl or C<sub>1-4</sub>alkyl-S(=O)<sub>n</sub>-; or Q is a radical of formula
- 15
- 20




wherein X and Y each independently are O, NR<sup>3</sup>, CH<sub>2</sub> or S, with R<sup>3</sup> being hydrogen or C<sub>1-4</sub>alkyl;

- 25 q is an integer with value 1 to 4;  
Z is O or NR<sup>4</sup> with R<sup>4</sup> being hydrogen or C<sub>1-4</sub>alkyl;  
r is an integer with value 1 to 3.

- A further interesting group comprises those compounds of formula (I) or (I') wherein Q is phenyl, naphthyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxy; amino; mono- or di(C<sub>1-6</sub>alkyl)amino; C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>2-6</sub>alkynyl; C<sub>3-6</sub>cycloalkyl; C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, amino, mono- or di(C<sub>1-4</sub>alkyl)amino; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylthio; C<sub>1-</sub>
- 30

<sub>6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxy carbonyl; C<sub>1-6</sub>alkylcarbonylamino; arylC<sub>1-6</sub>alkyloxy; aryloxy; polyhaloC<sub>1-6</sub>alkyl; polyhaloC<sub>1-6</sub>alkyloxy; polyhaloC<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyl-S(=O)<sub>n</sub>- or R<sup>1</sup>HN-S(=O)<sub>n</sub>-; or Q is a radical of formula



- 5 wherein X and Y each independently are O, NR<sup>3</sup>, CH<sub>2</sub> or S, with R<sup>3</sup> being hydrogen or C<sub>1-4</sub>alkyl;  
 q is an integer with value 1 to 4;  
 Z is O or NR<sup>4</sup> with R<sup>4</sup> being hydrogen or C<sub>1-4</sub>alkyl;  
 r is an integer with value 1 to 3.

10

Another interesting group comprises those compounds of formula (I) or (I') wherein Q is 3-pyridyl, 4-pyridyl, naphthalenyl, C<sub>3-6</sub>cycloalkyl, phenyl, 1,3-benzodioxolyl, 2,3-dihydro-benzofuranyl, 2,3-dihydro-1,4-benzodioxinyl, benzthiazolyl, indazolyl, benzimidazolyl or imidazopyridyl.

15

Also particular compounds are those compounds of formula (I) or (I') wherein Q is phenyl, 3-pyridyl, 4-pyridyl, benzthiazolyl or imidazopyridyl, in particular phenyl, each of said rings being optionally substituted with up to three substituents selected from halo, cyano, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy or polyhaloC<sub>1-6</sub>alkyl.

20

Each of the above-mentioned interesting groups of compounds of formula (I) or (I') describing a particular definition of L may be combined with each of the above-mentioned interesting groups of compounds of formula (I) or (I') describing a particular definition of Q.

25

Preferred compounds are selected from the group consisting of


- 2-thiazolamine, 4-imidazo[1,2-a]pyridin-3-yl-N-[3-(trifluoromethyl)phenyl];  
 2-thiazolamine, 4-imidazo[1,2-a]pyridin-3-yl-N-[4-(trifluoromethyl)phenyl];  
 2-thiazolamine, 4-(3-pyridinyl)-N-[3-(trifluoromethyl)phenyl];  
 30 2-thiazolamine, N-(3-chlorophenyl)-4-imidazo[1,2-a]pyridin-3-yl;  
 2-thiazolamine, 4-imidazo[1,2-a]pyridin-3-yl-N-(3-methylphenyl);  
 2-thiazolamine, 4-imidazo[1,2-a]pyridin-3-yl-N-[3-(methylthio)phenyl];  
 2-thiazolamine, N-(4-chlorophenyl)-4-imidazo[1,2-a]pyridin-3-yl;  
 2-thiazolamine, N-(3-bromophenyl)-4-imidazo[1,2-a]pyridin-3-yl;

- 2-thiazolamine, N-(2,3-dichlorophenyl)-4-imidazo[1,2-a]pyridin-3-yl;  
 2-thiazolamine, N-(2,3-dichlorophenyl)-4-(1H-pyrrolo[2,3-b]pyridin-3-yl);  
 2-thiazolamine, N-(4-bromophenyl)-4-imidazo[1,2-a]pyridin-3-yl;  
 2-thiazolamine, N-(2,3-dichlorophenyl)-4-imidazo[1,5-a]pyridin-3-yl;  
 5 2-thiazolamine, 4-imidazo[2,1-b]thiazol-5-yl-N-[3-(trifluoromethyl)phenyl];  
 2-thiazolamine, N-(2,3-dichlorophenyl)-4-imidazo[2,1-b]thiazol-5-yl;  
 2-thiazolamine, 4-(3-pyridinyl)-N-(3-methyl-4-fluorophenyl);  
 2-thiazolamine, 4-imidazo[1,2-a]pyridin-3-yl-N-(3-methyl-4-fluorophenyl);  
 the *N*-oxides, pharmaceutically acceptable addition salts, quaternary amines and  
 10 stereochemically isomeric forms thereof.


Also preferred compounds are selected from the group consisting of

- 2-thiazolamine, 4-(3-fluorophenyl)-*N*-phenyl;  
 2-thiazolamine, 4-(3-fluorophenyl)-*N*-[4-methoxyphenyl];  
 15 2-thiazolamine, 4-(3-fluorophenyl)-*N*-[4-(trifluoromethyl)phenyl]; and  
 2-thiazolamine, 4-(3-fluorophenyl)-*N*-[3-pyridyl];  
 the *N*-oxides, pharmaceutically acceptable addition salts, quaternary amines and  
 stereochemically isomeric forms thereof.


- 20 In general, the compounds of formula (I) may be prepared by reacting an intermediate of formula (II) or formula (III) or by reacting an intermediate of formula (II) and (III), wherein  $W_1$  represents a suitable leaving group, such as a halo atom, e.g. chloro or bromo, with an intermediate of formula (IV) in a suitable reaction-inert solvent, such as an alcohol, e.g. ethanol, or *N,N*-dimethylformamide.



- Compounds of formula (I), wherein L is substituted with amino, said L being represented by  $\text{NH}_2\text{-L}_1$ , and said compounds by formula (I-a), may be prepared by reacting an intermediate of formula (II), wherein Het is substituted with  $\text{C}_{1-6}\text{alkyl-C(=O)-NH-}$ , said Het being represented by  $\text{C}_{1-6}\text{alkyl-C(=O)-NH-Het}_1$ , and said intermediate being represented by formula (II-a), with an intermediate of formula (IV) in the presence of a suitable acid, such as for example hydrobromic acid and the like, in the presence of a suitable solvent, such as an alcohol, e.g. ethanol and the like, and water.



- Compounds of formula (I), wherein Q is substituted with amino, said Q being represented by  $\text{Q}_1\text{-NH}_2$ , and said compounds by formula (I-b), may be prepared by reducing an intermediate of formula (I-b-interm.), wherein Q is substituted with nitro, said Q being represented by  $\text{Q}_1\text{-NO}_2$ , in the presence of a suitable reducing agent, e.g. hydrogen, optionally in the presence of a suitable catalyst, e.g. palladium-on-charcoal, and a suitable catalyst poison, e.g. a thiophene solution. A suitable solvent for the above reaction is a reaction-inert solvent, for example, an alcohol, e.g. methanol.



Compounds of formula (I) may be converted into each other following art-known functional group transformation reactions, comprising those described hereinafter.


20

- The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboxylic acid or halo substituted benzenecarboxylic acid, e.g. 3-chlorobenzenecarboxylic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g.


25

t.butyl hydro-peroxide. Suitable solvents are, for example, water, lower alcohols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

- 5 Compounds of formula (I), wherein Q is substituted with cyano, said Q being represented by  $Q_1-CN$ , and said compounds by formula (I-c), may be converted into a compound of formula (I), wherein Q is substituted with carboxy, said Q being represented by  $Q_1-COOH$ , and said compound by formula (I-d), by reaction with a suitable acid, such as concentrated hydrochloric acid, in the presence of a suitable reaction-inert solvent, e.g. water.




- Compounds of formula (I), wherein L is substituted with  $C_{1-6}\text{alkyl-C(=O)-NH-}$ , said Het being represented by  $C_{1-6}\text{alkyl-C(=O)-NH-Het}_1$ , and said compounds being represented by formula (I-e), may be converted into a compound of formula (I-a), by reaction with a suitable acid, such as for example hydrobromic acid and the like, in the presence of a suitable solvent, such as water.




- In the following paragraphs, there are described several methods of preparing the intermediates in the foregoing preparations. A number of intermediates and starting materials are commercially available or are known compounds which may be prepared according to conventional reaction procedures generally known in the art.


- Intermediates of formula (II) can be prepared by reacting an intermediate of formula (V) with a suitable leaving group introducing agent of formula (VI), wherein  $W_1$  represents the leaving group and R represents the remaining of the agent, such as for example  $W_1-R$  representing  $Br_2$ , in the presence of a suitable solvent, such as a HBr solution, dioxane, acetic acid and the like.




- Alternatively, intermediates of formula (II) may also be prepared by Friedel-Crafts acylation in the presence of a suitable Lewis acid, for example by reacting an intermediate of formula (VII) with an intermediate of formula (VIII), wherein  $W_1$  and  $W_2$  represent a suitable leaving group, such as a halo atom, e.g. chloro, in the presence of  $AlCl_3$  and in the presence of a suitable solvent, e.g. carbon disulfide.




- Intermediates of formula (II) may also be prepared by acylating an intermediate of formula (VII-a), i.e. L having an acidic hydrogen atom, with an intermediate of formula (IX), with  $W_1$  as defined hereinabove, in the presence of a suitable base, e.g. lithium diisopropylamide, and a suitable reaction-inert solvent, e.g. tetrahydrofuran.



Intermediates of formula (II) may also be prepared by reacting an intermediate of formula (XI), with  $W_1$  as defined hereinabove, with a suitable acid, such as a HBr solution, in the presence of a suitable solvent, e.g. water.




- Intermediates of formula (III) may be prepared according to the first reaction procedure described above to prepare an intermediate of formula (II), thus by reacting an intermediate of formula (V) with an intermediate of formula (VI) in the presence of a suitable solvent, e.g. acetic acid, hydrobromic acid or the like.




- Intermediates of formula (V) may be prepared by reacting an intermediate of formula (XII), wherein  $W_3$  is a suitable leaving group, such as a halo atom, e.g. chloro, with an intermediate of formula (XIII) in the presence of *N,N*-dimethyl-4-pyridinamine and a suitable solvent, such as dichloromethane.

-31-



Intermediates of formula (XII), wherein  $\text{W}_3$  represents chloro, said intermediates being represented by formula (XII-a), can be prepared by reacting an intermediate of formula (XIV) with  $\text{SOCl}_2$ .




(XIV)

(XII-a)

5


Intermediates of formula (V), wherein L is Het and Het is an imidazo[1,2-a]pyrazinyl moiety as represented by formula (V-a), can be prepared by reacting an intermediate of formula (XV) with an intermediate of formula (XVI), wherein  $\text{W}_4$  is a suitable leaving group, such as a halo atom, e.g. bromo, in the presence of a suitable reaction-inert solvent, such as an alcohol, e.g. ethanol.

10



Intermediates of formula (V), wherein L is Het and Het is an imidazo[1,2-a]pyrimidinyl moiety as represented by formula (V-b), can be prepared by reacting an intermediate of formula (XVII) with an intermediate of formula (XVIII), wherein  $\text{W}_5$  represents a suitable leaving group, such as a halo atom, e.g. chloro, in the presence of a suitable reaction-inert solvent, such as methylene chloride.


15




Intermediates of formula (XVII) may be prepared by reacting an intermediate of formula (XIX) with an intermediate of formula (XX) in a reaction-inert solvent, such as toluene.

20


-32-




Intermediates of formula (IV) may be prepared by hydrolizing an intermediate of formula (XXI) in the presence of a suitable base, such as for example sodium hydroxide, and in the presence of a suitable solvent, such as an alcohol, e.g. ethanol and the like.




Intermediates of formula (XXI) may be prepared by reacting an intermediate of formula (XXII) with an intermediate of formula (XXIII) in the presence of a suitable solvent, such as tetrahydrofuran.



Intermediates of formula (XXII) may be prepared by hydrolyzing an intermediate of formula (XXIV) in the presence of a suitable acid, such as hydrobromic acid, hydrochloric acid, acetic acid and the like, or mixtures thereof, and in the presence of a suitable solvent, such as for example ethyl acetate.



Intermediates of formula (XXIV) may be prepared by reacting an intermediate of formula (XXV) with phosphorazidic acid diphenyl ester in the presence of a suitable base, such as *N,N*-diethyl-ethanamine, and in the presence of a suitable alcohol such as  $C_{1-6}$ alkylOH, e.g. ethanol, *t*-butanol and the like.



The compounds of the present invention show cytokine production modulating activity, in particular cytokine production inhibitory activity, more in particular proinflammatory cytokine production inhibitory activity. A cytokine is any secreted polypeptide that affects the function of other cells by modulating interactions between cells in the immune or inflammatory response. Examples of cytokines include

Interleukin-1 (IL-1) up to Interleukin-18 (IL-18), Tumor Necrosis Factor-alpha (TNF- $\alpha$ ), Tumor Necrosis Factor-beta (TNF- $\beta$ ). The present compounds also show inhibitory activity on the production of chemotactic cytokines or chemokines responsible for trafficking and activation of leucocytes. A chemokine production  
5 inhibited by the compounds of formula (I) or (I') is MCP-1 production (Monocyte Chemotactic Protein 1).

The cytokine production specifically inhibited by the compounds of formula (I) or (I') is TNF- $\alpha$  and/or Interleukin-12 (IL-12) production.

10

TNF- $\alpha$  is primarily produced by monocytes, macrophages, T and B lymphocytes, neutrophils, mast cells, tumour cells, fibroblasts, keratinocytes, astrocytes, microglial cells, smooth muscle cells and others. This proinflammatory cytokine is established at the pinnacle of proinflammatory cascades; it exerts a key role in the cytokine network  
15 with regard to the pathogenesis of many infectious, inflammatory and autoimmune diseases. Excessive or unregulated TNF- $\alpha$  production is implicated in mediating or exacerbating a number of diseases including rheumatoid arthritis, rheumatoid spondylitis, spondyloarthropathies, systemic lupus erythematosus, osteoarthritis, gouty arthritis, juvenile arthritis and other arthritic conditions, polychondritis, sclerodoma,  
20 Wegener granulomatosis, dermatomyositis, Steven-Johnson syndrome, idiopathic sprue, endocrine ophthalmopathy, Grave's disease, alveolitis, chronic hypersensitivity pneumonitis, primary biliary cirrhosis, uveitis, keratoconjunctivitis sicca and vernal keratoconjunctivitis, allergic rhinitis, pemphigus, eosinophilia, Loffler's syndrome, eosinophilic pneumonia, parasitic infestation, bronchopulmonary aspergillosis,  
25 polyarteritis nodosa, eosinophilic granuloma, eosinophil-related disorders affecting the airways occasioned by drug-reaction, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, cerebral malaria, adult respiratory distress syndrome, bronchitis (acute, arachidic, catarrhal, chronic, croupus, phthinoid bronchitis), chronic obstructive airway or pulmonary disease, pulmonary fibrosis,  
30 pneumoconiosis (aluminosis, anthracosis, asbestosis, chalicocis, ptilosis, siderosis, silicosis, tobaccosis, byssionosis), tuberculosis, silicosis, exacerbation of airways hyperreactivity to other drug therapy (e.g. aspirin or  $\beta$ -agonist therapy), pulmonary sarcoidosis, bone resorption diseases, meningitis, reperfusion injury, graft versus host reaction, allograft rejections, transplant rejections, fever and myalgias due to infection,  
35 such as influenza, cachexia (consequential to, e.g. bacterial, viral or parasitic, infection or to deprivation or deterioration of humoral or other organic function, or secondary to malignancy; malarial and vernal cachexia; cachexia resulting from dysfunction of the

pituitary, thyroid or thymus glands as well as uremic cachexia; cachexia secondary to acquired immune deficiency syndrome (AIDS)), AIDS, ARC (AIDS related complex), diabetes, cancer, angiogenesis, lymphoma, Kawasaki syndrome, Behçet's syndrome, aphthous ulceration, skin-related disorders such as psoriasis, eczema, burns, dermatitis, keloid formation, scar tissue formation, erythema nodosum leprosum, Crohn's disease, ulcerative colitis, inflammatory bowel disease, irritable bowel syndrome, pyresis, asthma (intrinsic, extrinsic, allergic, non-atopic, exercise induced and occupational and bacterial infection induced asthma), wheezy infant syndrome, multiple sclerosis, Parkinson's disease, pancreatitis, cardiac disease, congestive heart failure, myocardial infarction, acute liver failure, glomerulonephritis, therapy-associated syndromes comprising Jarisch-Herxheimer reaction, and syndromes associated with IL-2 infusion, anti-CD3 antibody infusion, hemodialysis, yellow fever vaccination.

TNF- $\alpha$  has also been shown to activate HIV (Human Immune deficiency Virus) replication in monocytes and/or macrophages. Therefore, inhibition of TNF- $\alpha$  production or activity aids in limiting HIV progression. TNF- $\alpha$  also plays a role in other viral infections, such as Hepatitis C, CMV (cytomegalovirus), influenza and herpes virus infections, including herpes simplex virus type-1, herpes simplex virus type-2, varicella-zoster virus, Epstein-Barr virus, human herpes virus-6,-7 and -8, pseudorabies and rhinotracheitis.

IL-12 is produced primarily by monocytes, macrophages and dendritic cells in response to bacteria, bacterial products (lipopolysaccharide) and immune signals. The production of IL-12 is regulated by other cytokines and endogenous mediators produced during inflammatory and immunological responses. IL-12 plays a central role in the immune system. Evidence obtained from animal models and human diseases suggests that inappropriate and protracted production of IL-12 and the ability of IL-12 to induce the generation of T helper 1 cell type responses may be instrumental in the development and maintenance of chronic inflammatory diseases, such as rheumatoid arthritis, collagen induced arthritis, allergic encephalitis, colitis, inflammatory bowel disease, Crohn's disease and multiple sclerosis, and in the triggering of autoimmune disorders, such as diabetes, or graft versus host diseases or shock. The adverse effects also include anemia (haemolytic, aplastic, pure red cell, idiopathic thrombocytopenia), neutropenia, lymphopenia, hepatosplenomegaly with mononuclear cell infiltration and pulmonary edema with interstitial cell infiltrates.

Excessive IL-12 production may accelerate the inflammatory progress of a disease, or the onset of the disease, such as rheumatoid arthritis, or it may also augment the disease severity.

Inhibition of TNF- $\alpha$  and/or IL-12 production by the compounds of formula (I) or (I') might offer an interesting, potentially less toxic alternative to non-specific immunosuppression (e.g. corticosteroids) in the treatment of chronic inflammatory and autoimmune diseases. The combined modulation of TNF- $\alpha$  and IL-12 production may ameliorate the treated disease to a greater extent than mono-therapy. The therapeutic effect of combining the suppression of both the immune and the inflammatory arm of a disease may provide additional clinical benefits. The present compounds are also indicated for use as co-therapeutic agents for use in conjunction with immunosuppressive and/or anti-inflammatory drugs, e.g. as potentiators of the therapeutic activity of said drugs, to reduce required dosaging or thus also potential side effects of said drugs. Immunosuppressive and/or anti-inflammatory drugs include for example cyclopeptide, cyclopeptolide or macrolide immunosuppressive or anti-inflammatory drugs, such as drugs belonging to the cyclosporin class, e.g. cyclosporine A or G, tacrolimus substances, ascomycin, rapamycin, glucocorticosteroid drugs, e.g. budesonide, beclamethasone, fluticasone, mometasone.

The compounds of formula (I) or (I') are useful in preventing or treating cytokine mediated diseases, and as such, inhibit, suppress or antagonize the production or activity of proinflammatory cytokines, such as TNF- $\alpha$  and/or IL-12.

Disorders mediated through TNF- $\alpha$  and/or IL-12 refers to any and all disorders and disease states in which TNF- $\alpha$  and/or IL-12 play a role, either by the cytokine itself, or by the cytokine causing another cytokine, such as for example IL-1 or IL-6, or a certain mediator to be released.

Due to their cytokine production inhibitory activity, in particular their proinflammatory cytokine production inhibitory activity, more in particular their TNF- $\alpha$  and/or IL-12 inhibitory activity, the compounds of formula (I) or (I'), their N-oxides, pharmaceutically acceptable addition salts, quaternary amines and stereochemically isomeric forms are useful in the treatment or prevention of diseases or conditions mediated through cytokines, in particular diseases or conditions related to excessive or unregulated production of proinflammatory cytokines, such as TNF- $\alpha$  and/or IL-12, comprising inflammatory diseases or auto-immune diseases. Diseases or conditions related to an excessive or unregulated production of proinflammatory cytokines comprise rheumatoid arthritis, rheumatoid spondylitis, spondyloarthropathies, systemic lupus erythematosus, osteoarthritis, gouty arthritis, juvenile arthritis and other arthritic

- conditions, polychondritis, sclerodoma, Wegener granulomatosis, dermatomyositis, Steven-Johnson syndrome, idiopathic sprue, endocrine ophthalmopathy, Graves' disease, alveolitis, chronic hypersensitivity pneumonitis, primary billiary cirrhosis, uveitis, keratoconjunctivitis sicca and vernal keratoconjunctivitis, allergic rhinitis, pemphigus,
- 5 eosinophilia, Loffler's syndrome, eosinophilic pneumonia, parasitic infestation, bronchopulmonary aspergillosis, polyarteritis nodosa, eosinophilic granuloma, eosinophil-related disorders affecting the airways occasioned by drug-reaction, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, cerebral malaria, adult respiratory distress syndrome, bronchitis (acute, arachidic, catarrhal,
- 10 chronic, croupus, phthinoid bronchitis), chronic obstructive airway or pulmonary disease, pulmonary fibrosis, tuberculosis, pneumoconiosis (aluminosis, anthracosis, asbestosis, chalicocis, ptilosis, siderosis, silicosis, tobaccosis, byssionosis), exacerbation of airways hyperreactivity to other drug therapy (e.g. aspirin or  $\beta$ -agonist therapy), silicosis, pulmonary sarcoidosis, bone resorption diseases, meningitis, allergic
- 15 encephalitis, reperfusion injury, graft versus host reaction, allograft rejections, transplant rejections, fever and myalgias due to infection, such as influenza, cachexia (consequential to, e.g. bacterial, viral or parasitic, infection or to deprivation or deterioration of humoral or other organic function, or secondary to malignancy; malarial and vernal cachexia; cachexia resulting from dysfunction of the pituitary,
- 20 thyroid or thymus glands as well as uremic cachexia; cachexia secondary to acquired immune deficiency syndrome (AIDS)), AIDS, ARC (AIDS related complex), diabetes, cancer, angiogenesis, lymphoma, Kawasaki syndrome, Behçet's syndrome, aphthous ulceration, skin-related disorders such as psoriasis, eczema, burns, dermatitis, keloid formation, scar tissue formation, erythema nodosum leprosum, Crohn's disease,
- 25 ulcerative colitis, inflammatory bowel disease, irritable bowel syndrome, pyresis, asthma (intrinsic, extrinsic, allergic, non-atopic, exercise induced and occupational and bacterial infection induced asthma), wheezy infant syndrome, multiple sclerosis, Parkinson's disease, pancreatitis, cardiac disease, congestive heart failure, myocardial infarction, acute liver failure, glomerulonephritis, therapy-associated syndromes
- 30 comprising Jarisch-Herxheimer reaction, and syndromes associated with IL-2 infusion, anti-CD3 antibody infusion, hemodialysis, yellow fever vaccination, HIV or other viral infections, such as Hepatitis C, CMV, influenza and herpes virus infections, pseudorabies and rhinotracheitis, angiofollicular lymphoid hyperplasia, anemia (haemolytic, aplastic, pure red cell, idiopathic thrombocytopenia), neutropenia,
- 35 lymphopenia, hepatosplenomegaly with mononuclear cell infiltration and pulmonary edema with interstitial cell infiltrates; or to prevent these diseases. In particular, the

compounds of formula (I) or (I') can be used to treat rheumatoid arthritis, Crohn's disease, irritable bowel disease or colitis.

5 The cytokine production inhibitory activity of the compounds of formula (I) or (I') such as the inhibition of TNF- $\alpha$  and/or IL-12 production, may be demonstrated in the *in vitro* test "Inhibition of cytokine production in human whole blood cultures". Suitable *in vivo* tests are "Determination of cytokine in serum of LPS (lipopolysaccharide) and anti-CD3 challenged mice", "Inhibition of LPS-galactosamine induced shock in mice", "Inhibition of collagen induced arthritis in mice".

10

The compounds of formula (I) or (I') may also inhibit Interleukin-6 (IL-6).

The present compounds also have a selective affinity for adenosine A<sub>3</sub> receptors. Therefore, they can be used to prevent and/or treat adenosine related diseases such as  
15 asthma, allergosis, inflammation, Addison's disease, autoallergic hemolytic anemia, Crohn's disease, psoriasis, rheumatism, diabetes.

The present compounds may also act as intermediates for the preparation of further thiazolyl derivatives.

20

In view of the above described pharmacological properties, the compounds of formula (I) or (I') or any subgroup thereof, their N-oxides, pharmaceutically acceptable addition salts, quaternary amines and stereochemically isomeric forms, may be used as a  
25 medicine. In particular, the present compounds can be used for the manufacture of a medicament for treating or preventing diseases mediated through cytokines, more in particular diseases mediated through TNF- $\alpha$  and/or IL-12, such as inflammatory and auto-immune diseases. The present compounds can also be used for the manufacture of a medicament for treating or preventing diseases mediated through activation of the adenosine A<sub>3</sub> receptor.

30

In view of the utility of the compounds of formula (I) or (I'), there is provided a method of treating warm-blooded animals, including humans, suffering from or a method of preventing warm-blooded animals, including humans, to suffer from  
35 diseases mediated through cytokines, in particular mediated through TNF- $\alpha$  and/or IL-12, such as inflammatory and auto-immune diseases. There is also provided a method of treating warm-blooded animals, including humans, suffering from or a method of preventing warm-blooded animals, including humans, to suffer from diseases mediated

through activation of the adenosine A<sub>3</sub> receptor. Said methods comprise the administration, preferably oral administration, of an effective amount of a compound of formula (I) or (I'), a *N*-oxide form, a pharmaceutically acceptable addition salt, a quaternary amine or a possible stereoisomeric form thereof, to warm-blooded animals, including humans.

The present invention also provides compositions for preventing or treating diseases mediated through cytokines or mediated through activation of the adenosine A<sub>3</sub> receptor comprising a therapeutically effective amount of a compound of formula (I) and a pharmaceutically acceptable carrier or diluent.

The compounds of the present invention or any subgroup thereof may be formulated into various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, optionally in addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, particularly, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches, sugars, kaolin, diluents, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a

suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be  
5 administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment. The compounds of the present invention may also be administered via inhalation or insufflation by means of methods and formulations employed in the art for administration via this way. Thus, in general the compounds of the present invention may be administered to the lungs in the form of a solution, a suspension or a dry  
10 powder. Any system developed for the delivery of solutions, suspensions or dry powders via oral or nasal inhalation or insufflation are suitable for the administration of the present compounds.

To aid solubility of the compounds of formula (I), suitable ingredients, e.g.  
15 cyclodextrins, may be included in the compositions. Appropriate cyclodextrins are  $\alpha$ -,  $\beta$ -,  $\gamma$ -cyclodextrins or ethers and mixed ethers thereof wherein one or more of the hydroxy groups of the anhydroglucose units of the cyclodextrin are substituted with  $C_{1-6}$ alkyl, particularly methyl, ethyl or isopropyl, e.g. randomly methylated  $\beta$ -CD; hydroxy $C_{1-6}$ alkyl, particularly hydroxyethyl, hydroxy-propyl or hydroxybutyl;  
20 carboxy $C_{1-6}$ alkyl, particularly carboxymethyl or carboxy-ethyl;  $C_{1-6}$ alkylcarbonyl, particularly acetyl. Especially noteworthy as complexants and/or solubilizers are  $\beta$ -CD, randomly methylated  $\beta$ -CD, 2,6-dimethyl- $\beta$ -CD, 2-hydroxyethyl- $\beta$ -CD, 2-hydroxyethyl- $\gamma$ -CD, 2-hydroxypropyl- $\gamma$ -CD and (2-carboxymethoxy)propyl- $\beta$ -CD, and in particular 2-hydroxypropyl- $\beta$ -CD (2-HP- $\beta$ -CD).

25 The term mixed ether denotes cyclodextrin derivatives wherein at least two cyclodextrin hydroxy groups are etherified with different groups such as, for example, hydroxy-propyl and hydroxyethyl.

30 The average molar substitution (M.S.) is used as a measure of the average number of moles of alkoxy units per mole of anhydroglucose. The average substitution degree (D.S.) refers to the average number of substituted hydroxyls per anhydroglucose unit. The M.S. and D.S. value can be determined by various analytical techniques such as nuclear magnetic resonance (NMR), mass spectrometry (MS) and infrared  
35 spectroscopy (IR). Depending on the technique used, slightly different values may be obtained for one given cyclodextrin derivative. Preferably, as measured by mass spectrometry, the M.S. ranges from 0.125 to 10 and the D.S. ranges from 0.125 to 3.

Other suitable compositions for oral or rectal administration comprise particles consisting of a solid dispersion comprising a compound of formula (I) and one or more appropriate pharmaceutically acceptable water-soluble polymers.

5

The term "a solid dispersion" used hereinafter defines a system in a solid state (as opposed to a liquid or gaseous state) comprising at least two components, in casu the compound of formula (I) and the water-soluble polymer, wherein one component is dispersed more or less evenly throughout the other component or components ( in case additional pharmaceutically acceptable formulating agents, generally known in the art, are included, such as plasticizers, preservatives and the like). When said dispersion of the components is such that the system is chemically and physically uniform or homogenous throughout or consists of one phase as defined in thermo-dynamics, such a solid dispersion will be called "a solid solution". Solid solutions are preferred physical systems because the components therein are usually readily bioavailable to the organisms to which they are administered. This advantage can probably be explained by the ease with which said solid solutions can form liquid solutions when contacted with a liquid medium such as the gastro-intestinal juices. The ease of dissolution may be attributed at least in part to the fact that the energy required for dissolution of the components from a solid solution is less than that required for the dissolution of components from a crystalline or microcrystalline solid phase.

The term "a solid dispersion" also comprises dispersions which are less homogenous throughout than solid solutions. Such dispersions are not chemically and physically uniform throughout or comprise more than one phase. For example, the term "a solid dispersion" also relates to a system having domains or small regions wherein amorphous, microcrystalline or crystalline compound of formula (I), or amorphous, microcrystalline or crystalline water-soluble polymer, or both, are dispersed more or less evenly in another phase comprising water-soluble polymer, or compound of formula (I), or a solid solution comprising compound of formula (I) and water-soluble polymer. Said domains are regions within the solid dispersion distinctively marked by some physical feature, small in size, and evenly and randomly distributed throughout the solid dispersion.

Various techniques exist for preparing solid dispersions including melt-extrusion, spray-drying and solution-evaporation.

The solution-evaporation process comprises the following steps :

- a) dissolving the compound of formula (I) and the water-soluble polymer in an appropriate solvent, optionally at elevated temperatures;
- b) heating the solution resulting under point a), optionally under vacuum, until the solvent is evaporated. The solution may also be poured onto a large surface so as to form a thin film, and evaporating the solvent therefrom.

In the spray-drying technique, the two components are also dissolved in an appropriate solvent and the resulting solution is then sprayed through the nozzle of a spray dryer followed by evaporating the solvent from the resulting droplets at elevated temperatures.

The preferred technique for preparing solid dispersions is the melt-extrusion process comprising the following steps :

- a) mixing a compound of formula (I) and an appropriate water-soluble polymer,
- b) optionally blending additives with the thus obtained mixture,
- c) heating and compounding the thus obtained blend until one obtains a homogenous melt,
- d) forcing the thus obtained melt through one or more nozzles; and
- e) cooling the melt till it solidifies.

The terms "melt" and "melting" should be interpreted broadly. These terms not only mean the alteration from a solid state to a liquid state, but can also refer to a transition to a glassy state or a rubbery state, and in which it is possible for one component of the mixture to get embedded more or less homogeneously into the other. In particular cases, one component will melt and the other component(s) will dissolve in the melt thus forming a solution, which upon cooling may form a solid solution having advantageous dissolution properties.

After preparing the solid dispersions as described hereinabove, the obtained products can be optionally milled and sieved.

The solid dispersion product may be milled or ground to particles having a particle size of less than 600  $\mu\text{m}$ , preferably less than 400  $\mu\text{m}$  and most preferably less than 125  $\mu\text{m}$ .

The particles prepared as described hereinabove can then be formulated by conventional techniques into pharmaceutical dosage forms such as tablets and capsules.

It will be appreciated that a person of skill in the art will be able to optimize the parameters of the solid dispersion preparation techniques described above, such as the most appropriate solvent, the working temperature, the kind of apparatus being used,  
5 the rate of spray-drying, the throughput rate in the melt-extruder

The water-soluble polymers in the particles are polymers that have an apparent viscosity, when dissolved at 20°C in an aqueous solution at 2 % (w/v), of 1 to 5000 mPa.s more preferably of 1 to 700 mPa.s, and most preferred of 1 to 100 mPa.s. For  
10 example, suitable water-soluble polymers include alkylcelluloses, hydroxyalkyl-celluloses, hydroxyalkyl alkylcelluloses, carboxyalkylcelluloses, alkali metal salts of carboxyalkylcelluloses, carboxyalkylalkylcelluloses, carboxyalkylcellulose esters, starches, pectines, chitin derivatives, di-, oligo- and polysaccharides such as trehalose, alginic acid or alkali metal and ammonium salts thereof, carrageenans, galactomannans,  
15 tragacanth, agar-agar, gummi arabicum, guar gummi and xanthan gummi, polyacrylic acids and the salts thereof, polymethacrylic acids and the salts thereof, methacrylate copolymers, polyvinylalcohol, polyvinylpyrrolidone, copolymers of polyvinylpyrrolidone with vinyl acetate, combinations of polyvinylalcohol and polyvinylpyrrolidone, polyalkylene oxides and copolymers of ethylene oxide and  
20 propylene oxide. Preferred water-soluble polymers are hydroxypropyl methylcelluloses.

Also one or more cyclodextrins can be used as water soluble polymer in the preparation of the above-mentioned particles as is disclosed in WO 97/18839. Said cyclodextrins  
25 include the pharmaceutically acceptable unsubstituted and substituted cyclodextrins known in the art, more particularly  $\alpha$ ,  $\beta$  or  $\gamma$  cyclodextrins or the pharmaceutically acceptable derivatives thereof.

Substituted cyclodextrins which can be used to prepare the above described particles  
30 include polyethers described in U.S. Patent 3,459,731. Further substituted cyclodextrins are ethers wherein the hydrogen of one or more cyclodextrin hydroxy groups is replaced by C<sub>1-6</sub>alkyl, hydroxyC<sub>1-6</sub>alkyl, carboxy-C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxycarbonylC<sub>1-6</sub>alkyl or mixed ethers thereof. In particular such substituted cyclodextrins are ethers wherein the hydrogen of one or more cyclodextrin hydroxy  
35 groups is replaced by C<sub>1-3</sub>alkyl, hydroxyC<sub>2-4</sub>alkyl or carboxyC<sub>1-2</sub>alkyl or more in particular by methyl, ethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, carboxy-methyl or carboxyethyl.

Of particular utility are the  $\beta$ -cyclodextrin ethers, e.g. dimethyl- $\beta$ -cyclodextrin as described in Drugs of the Future, Vol. 9, No. 8, p. 577-578 by M. Nogradi (1984) and polyethers, e.g. hydroxypropyl  $\beta$ -cyclodextrin and hydroxyethyl  $\beta$ -cyclodextrin, being  
5 examples. Such an alkyl ether may be a methyl ether with a degree of substitution of about 0.125 to 3, e.g. about 0.3 to 2. Such a hydroxypropyl cyclodextrin may for example be formed from the reaction between  $\beta$ -cyclodextrin and propylene oxide and may have a MS value of about 0.125 to 10, e.g. about 0.3 to 3.

10 Another type of substituted cyclodextrins is sulfobutylcyclodextrines.

The ratio of the compound of formula (I) over the water soluble polymer may vary widely. For example ratios of 1/100 to 100/1 may be applied. Interesting ratios of the compound of formula (I) over cyclodextrin range from about 1/10 to 10/1. More  
15 interesting ratios range from about 1/5 to 5/1.

It may further be convenient to formulate the compounds of formula (I) in the form of nanoparticles which have a surface modifier adsorbed on the surface thereof in an amount sufficient to maintain an effective average particle size of less than 1000 nm.  
20 Useful surface modifiers are believed to include those which physically adhere to the surface of the compound of formula (I) but do not chemically bond to said compound.

Suitable surface modifiers can preferably be selected from known organic and inorganic pharmaceutical excipients. Such excipients include various polymers, low molecular  
25 weight oligomers, natural products and surfactants. Preferred surface modifiers include nonionic and anionic surfactants.

Yet another interesting way of formulating the compounds of formula (I) involves a pharmaceutical composition whereby the compounds of formula (I) are incorporated in  
30 hydrophilic polymers and applying this mixture as a coat film over many small beads, thus yielding a composition which can conveniently be manufactured and which is suitable for preparing pharmaceutical dosage forms for oral administration.

Said beads comprise a central, rounded or spherical core, a coating film of a  
35 hydrophilic polymer and a compound of formula (I) and optionally a seal-coating layer.

Materials suitable for use as cores in the beads are manifold, provided that said materials are pharmaceutically acceptable and have appropriate dimensions and firmness. Examples of such materials are polymers, inorganic substances, organic substances, and saccharides and derivatives thereof.

5

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, suppositories, injectable solutions or suspensions and the like, and segregated multiples thereof.

10

15 The present compounds are orally active compounds, and are preferably orally administered.

The exact dosage and frequency of administration depends on the particular compound of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight, sex, extent of disorder and general physical condition of the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention.

20

25

The compounds of formula (I) may also be used in combination with other conventional anti-inflammatory or immunosuppressive agents, such as steroids, cyclooxygenase-2 inhibitors, non-steroidal anti-inflammatory drugs, TNF- $\alpha$  antibodies, such as for example acetyl salicylic acid, bufexamac, diclofenac potassium, sulindac, diclofenac sodium, ketorolac trometamol, tolmetine, ibuprofen, naproxen, naproxen sodium, tiaprofen acid, flurbiprofen, mefenamic acid, niflumonic acid, meclofenamate, indomethacin, proglumetacine, ketoprofen, nabumetone, paracetamol, piroxicam, tenoxicam, nimesulide, fentanylbutazon, tramadol, beclomethasone dipropionate, betamethasone, beclomethasone, budesonide, fluticasone, mometasone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, triamcinolone, celecoxib, rofecoxib, infliximab, leflunomide, etanercept, CPH 82, methotrexate, sulfasalazine, antilymphocytory immunoglobulines, antithymocytory

30

35

immunoglobulines, azathioprine, cyclosporine, tacrolimus substances, ascomycin, rapamycin, muromonab-CD3.

Thus, the present invention also relates to the combination of a compound of formula (I) and another anti-inflammatory or immunosuppressive agent. Said combination may be used as a medicine. The present invention also relates to a product containing (a) a compound of formula (I), and (b) another anti-inflammatory or immunosuppressive compound, as a combined preparation for simultaneous, separate or sequential use in the treatment of diseases related to an excessive or unregulated cytokine production. The different drugs may be combined in a single preparation together with pharmaceutically acceptable carriers.

### Experimental part

#### A. Preparation of the intermediate compounds

##### Example A1

2-Bromo-acetoacetaldehyde (0.1 mol) was added portionwise to pyrazinamine (0.1 mol) in ethanol (200 ml) while stirring. The reaction mixture was stirred and refluxed for one hour, then allowed to cool to room temperature. The precipitate was filtered off and dried. Yield: 13.5 g of 1-(imidazo[1,2-a]pyrazin-3-yl)ethanone (55%) (interm. 1).

##### Example A2

a) A mixture of 2-pyrimidinamine (0.5 mol) and 1,1-dimethoxy-*N,N*-dimethylmethanamine (0.55 mol) in methylbenzene (500 ml) was stirred and refluxed for 2 hours. The reaction mixture was cooled and the solvent was evaporated. Yield:  $\pm$  75 g of *N,N*-dimethyl-*N'*-(2-pyrimidinyl)methanimidamide (interm. 2). b) A mixture of intermediate (2) (0.066 mol) and 1-chloro-2-propanone (0.13 mol) in  $\text{CH}_2\text{Cl}_2$  (500 ml) was stirred and refluxed for 48 hours. The reaction mixture was cooled and the solvent was evaporated. The residue was crystallized from  $\text{CH}_3\text{CN}$ , filtered off, washed and dried. Yield: 6.9 g of imidazo[1,2-a]pyrimidin-3-ylethanone (65.1%) (interm. 3).

##### Example A3

a) A mixture of 6-(trifluoromethyl)-3-pyridinecarboxylic acid (0.026 mol) in thionyl chloride (50 ml) was stirred and refluxed for 2 hours. The solvent was evaporated. Yield : 5.2g of 6-(trifluoromethyl)-3-pyridinecarbonyl chloride (interm. 4)  
b) A mixture of 2,2-dimethyl-1,3-dioxane-4,6-dione (0.025 mol) in dichloromethane (150ml) was stirred under  $\text{N}_2$  flow and cooled to  $0^\circ\text{C}$ . *N,N*-Dimethyl-4-pyridinamine (0.055 mol) was dissolved in dichloromethane (50 ml) and added dropwise to the first solution at  $0^\circ\text{C}$ . This reaction mixture was stirred for 30 minutes without an ice-bath. The mixture was again cooled and intermediate 4 (0.025 mol) was dissolved in

dichloromethane (100 ml) and added dropwise to the first solution at 0°C. The reaction mixture was stirred for 2 hours at 0°C and overnight at room temperature under N<sub>2</sub> flow. The solvent was evaporated and the residue was taken up in ethyl acetate and washed with HCl 1N (30 ml) and H<sub>2</sub>O (70 ml) and again with H<sub>2</sub>O (2x). The separated  
5 organic layer was dried, filtered and the solvent was evaporated. Yield : 6.1g of 1-[6-(trifluoromethyl)-3-pyridinyl]ethanone ( interm. 5)

#### Example A4

Reference method: Lipinski et al. J.Org.Chem. 1984,49,50. A solution of acetyl chloride (0.072 mol) in dichloromethane (10 ml) was added dropwise to a mixture of 1-  
10 (2-methyl-1*H*-imidazol-4-yl)ethanone (0.024 mol) and *N,N*-diethylethanamine (0.072 mol) in dichloromethane (230 ml). The mixture was stirred for 1 hour. *N,N*-diethylethanamine (0.75 g) was added again. The mixture was washed very shortly with ice water (50 ml) and separated into its layers. The aqueous layer was extracted twice with CH<sub>2</sub>Cl<sub>2</sub> (30 ml). The combined organic layer was dried (MgSO<sub>4</sub>), filtered  
15 and the solvent was evaporated. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (100 ml). trimethyloxonium tetrafluoroborate (0.053 mol) was added. Na<sub>2</sub>CO<sub>3</sub> (80 ml) was added. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0 to 98/2). The pure fractions were collected and the  
20 solvent was evaporated. Yield: 3.4g of 1-(1,2-dimethyl-1*H*-imidazol-5-yl)ethanone (interm. 6)

#### Example A5

a) 6-Chloro-imidazo[1,2-*a*]pyridine (0.1 mol) was dissolved in CS<sub>2</sub> (400 ml). The solution was warmed. AlCl<sub>3</sub> (0.3 mol) was added portionwise (exothermic temperature  
25 rise to reflux temperature). A solution of chloroacetyl chloride (0.2 mol) in CS<sub>2</sub> (100 ml) was added dropwise and the reaction mixture was stirred and refluxed for 4 hours, then stirred overnight at room temperature. The mixture was decomposed with ice (200 g). CH<sub>3</sub>OH (100 ml) was added. 1N HCl (100 ml) was added and the mixture was stirred for 2 hours. The precipitate was filtered off, rinsed with 2-propanone and  
30 dried. Yield: 8.86 g of 2-chloro-1-(6-chloroimidazo[1,2-*a*]pyridin-3-yl)ethanone monohydrochloride (interm. 7). The filtrate was alkalized with Na<sub>2</sub>CO<sub>3</sub>, then with 50% NaOH. This mixture was extracted with ethyl acetate (3 x). The separated organic layer was dried, filtered and the solvent evaporated. The residue was dissolved in 2-propanone and converted into the hydrochloric acid salt (1:1) with HCl/2-propanol.  
35 The precipitate was filtered off and dried. Yield : 1.81g of intermediate (7). Total yield : 10.67g (40.2%) of intermediate (7).

- b) Reaction under N<sub>2</sub> atmosphere. Tetrahydrofuran (700 ml) was cooled to -70°C. n-Butyllithium 2.5M in hexane (100 ml) was added. A solution of *N*-(1-methylethyl)-2-propanamine (0.22 mol) in tetrahydrofuran (100 ml) was added dropwise at -70°C, then warmed slowly to -40°C and stirred for 30 minutes at -40°C. The reaction mixture was re-cooled to -70°C. A solution of imidazo[1,5-a]pyridine (0.2 mol) in tetrahydrofuran (100 ml) was added dropwise and the reaction mixture was stirred for 2 hours, allowing the temperature to rise to  $\pm$  -30°C. The reaction mixture was re-cooled to -70°C. A solution of *N,N*-dimethyl-2-chloroacetamide (0.22 mol) in tetrahydrofuran (100 ml) was added dropwise. The cooling bath was removed and the reaction mixture was stirred until the temperature reached  $\pm$  0°C. The reaction mixture was cooled, decomposed with ice and 2N HCl. The layers were separated. The water layer was extracted twice with ethyl acetate. The separated organic layer was dried, filtered and the solvent evaporated. Yield: 24 g of 2-chloro-1-(imidazo[1,5-a]pyridin-3-yl)ethanone (62%) (interm. 8).
- c) Intermediate (1) (0.02 mol) in HBr 48% (90 ml) was stirred at 70°C. A solution of Br<sub>2</sub> (0.02 mol) in HBr 48% (10 ml) was added dropwise and the reaction mixture was stirred for one hour at 70°C. The solvent was evaporated. The residue was stirred in 2-propanone with a small amount of ethanol, filtered off and dried. Yield: 6.15 g of 2-bromo-1-(imidazo[1,2-a]pyrazin-3-yl)ethanone monohydrobromide (interm. 9).
- d) 1-(1*H*-indazol-3-yl)ethanone (0.01 mol) was stirred in 1,4-dioxane (100 ml), at room temperature. A solution of Br<sub>2</sub> (0.01 mol) in 1,4-dioxane (20 ml) was added dropwise and the resulting reaction mixture was stirred overnight at room temperature. The precipitate was filtered off and the filtrate was evaporated. The residue was crystallized from CH<sub>3</sub>OH, filtered off and dried. Yield: 0.73 g of 2-bromo-1-(1*H*-indazol-3-yl)ethanone (interm. 10).
- e) Intermediate (3) (0.15 mol) was dissolved in acetic acid (250 ml). A solution of Br<sub>2</sub> (0.3 mol) in acetic acid (40 ml) was added dropwise at room temperature and the resulting reaction mixture was stirred for 2 hours at 100°C (steam bath). The reaction mixture was cooled to 0°C, then stirred overnight at room temperature. The precipitate was filtered off, washed and dried (in vacuo). Yield: 40.4 g (84.2%, mixture of two major compounds). HPLC separation gave two fraction groups. The solvent of each group was evaporated. Yield: 17 g of 2,2-dibromo-1-(imidazo[1,2-a]pyrimidin-3-yl)ethanone (interm. 8) and 7.2 g of 2-bromo-1-(imidazo[1,2-a]pyrimidin-3-yl)ethanone monohydrobromide (interm. 11).
- f) 1-(6-chloroimidazo[1,2-b]pyridazin-3-yl)ethanone (0.005 mol) was dissolved in a solution of hydrobromide 48% (15 ml). The mixture was heated to  $\pm$  70 °C. Br<sub>2</sub> (0.005 mol) was added dropwise over 15 minutes. The reaction mixture was stirred overnight


at room temperature. The precipitate was filtered off, washed, then suspended in 2-propanone. The precipitate was filtered off, washed and dried. Yield: 1.2 g of 2,2-dibromo-1-(6-chloroimidazo[1,2-b]pyridazin-3-yl)ethanone (interm. 12).

#### Example A6


- 5 a) *N,N*-diethylethanamine (2.61 g) was added to a mixture of 6-(trifluoromethyl)-3-pyridinecarboxylic acid (0.025 mol) in *t*-butanol (100 ml). The mixture was warmed up to 90°C. Phosphorazidic acid, diphenyl ester (0.025 mol) was added dropwise ( $N_2$ -development). The mixture was stirred at 90°C overnight. The solvent was evaporated. The residue (16.98 g) was purified by column chromatography over silica  
10 gel (eluent  $CH_2Cl_2$  100%). The pure fractions were collected and the solvent was evaporated. Yield: 6.3 g (96%) of carbamic acid, (6-trifluoromethyl-3-pyridinyl), 1,1-dimethylethyl ester (interm. 13)
- b) HBr/acetic acid (30 ml) was added to a mixture of intermediate 13 (0.02 mol) and ethyl acetate (150 ml) (a precipitate was formed immediately). EtOH was added. More  
15 HBr/acetic acid (10 ml) was added. The solvent was evaporated. The residue was taken up in ethyl acetate. NaOH (1M) was added. The mixture was extracted. The organic layer was separated, dried, filtered and the solvent was evaporated. HCl 1M (100 ml) was added. The solution was stirred at 80°C for 4 hours. The solvent was evaporated. NaOH (1M) was added. The mixture was extracted with  $CH_2Cl_2$  (3X 100 ml). The  
20 combined organic layer was dried, filtered and the solvent was evaporated. Yield: 2.64 g of 6-(trifluoromethyl)-3-pyridinamine (interm. 14).
- c) A solution of benzoyl isothiocyanate (0.016 mol) in tetrahydrofuran (50 ml) was added at room temperature to a mixture of intermediate 14 (0.016 mol) in tetrahydrofuran (200 ml). The mixture was stirred overnight. The solvent was  
25 evaporated. The residue was stirred in diisopropyl ether. The precipitate was filtered off and dried in vacuo at 40°C. Yield: 3.189 g (61.3%) of *N*-[[6-(trifluoromethyl)-3-pyridinyl-amino]thioxomethyl]benzamide (interm. 15).
- d) A mixture of intermediate 15 (0.0098 mol) and NaOH 1M (0.01 mol) in ethanol (150 ml) was stirred and refluxed for 30 minutes and then cooled.  $MgSO_4$  was added. The  
30 mixture was filtered and the filtrate was evaporated. The residue was stirred in diisopropyl ether, stirred and refluxed, cooled, filtered and dried. Yield: 1.178 g (54.3%) of [6-(trifluoromethyl)-3-pyridinyl] thiourea (interm. 16).

The following intermediates were prepared analogous to one of the above examples  
35 (the example number according to which they were prepared is indicated between square brackets after the intermediate number).


-49-



interm. 17 [A5d]




5 interm. 18 [A5c]




hydrobromide (1:1);interm. 19 [A5c]

10



hydrochloride (1:1);interm. 20 [A5a]



15 hydrobromide (1:1);interm. 21 [A1]

### B. Preparation of the final compounds

#### Example B1

- a) A mixture of 2-chloro-1-(imidazo[2,1-b]thiazol-5-yl)ethanone monohydrochloride (0.0025 mol), prepared according to A5a), and intermediate 16 (0.0025 mol) in ethanol (50 ml) was stirred at 80°C for 10 hours and then cooled. The precipitate was filtered off and dried. Yield: 0.54g of 4-(imidazo[2,1-b]thiazol-5-yl)-N-[(6-trifluoromethyl)-3-pyridinyl]-2-thiazolamine monohydrochloride ; mp 242°C (comp. 568).
- b) A mixture of intermediate (10) (0.001 mol) and (4-chlorophenyl)thiourea (0.001 mol) in ethanol (10 ml) was stirred for 3 hours at  $\pm 70^\circ\text{C}$ , then stirred overnight at room temperature. The precipitate was filtered off and dried. Yield: 0.33 g of N-(4-chlorophenyl)-4-imidazo[1,2-a]pyrazin-3-yl-2-thiazolamine monohydrobromide (comp. 2).
- c) A mixture of intermediate (11) (0.005 mol) and 3-pyridinylthiourea (0.005 mol) in ethanol (50 ml) was stirred and refluxed for 12 hours, then cooled and the resulting precipitate was filtered off, washed and dried (vacuum). Yield: 0.2 g of N-(4-imidazo[1,2-a]pyrimidin-3-yl-2-thiazolyl)-3-pyridinamine monohydrobromide (10.5%) (comp. 3).
- d) A mixture of 2-bromo 1-(5-methyl-3-pyridinyl)ethanone (0.00125 mol) and 2,2-dibromo 1-(5-methyl-3-pyridinyl)ethanone (0.00125 mol), both prepared according to A5e), and [3-(trifluoromethyl)-phenyl]thiourea in ethanol (25 ml) was stirred and refluxed for 3 hours. The reaction mixture was stirred overnight at room temperature.

A solid was formed, filtered off, washed and dried (vacuum). Yield : 0.4 g of N-[3-(trifluoromethyl)phenyl]-4-[5-methyl-3-pyridinyl]-2-thiazolamine monohydrobromide (comp. 626).

#### Example B2

- 5 A mixture of N-(3-nitro-phenyl-4-imidazo[1,2-a]pyridin-3-yl-2-thiazolamine, (0.003 mol), prepared according to the synthesis procedure described under B1a-2), in methanol (150 ml) was hydrogenated with palladium-on-charcoal 10% (1 g) as a catalyst in the presence of thiophene 4% in diisopropylether (1 ml). After uptake of hydrogen (3 equivalents), the catalyst was filtered off and the filtrate was evaporated.
- 10 The residue was dissolved in ethanol and converted into the hydrochloric acid salt (1:2) with HCl/2-propanol. The precipitate was filtered off and dried. Yield: 0.85 g of N-(4-imidazo[1,2-a]pyridin-3-yl-2-thiazolyl)-1,3-benzenediamine dihydrochloride monohydrate (comp 5).

#### Example B3

- 15 A mixture of compound (6) (see Table 2) (0.0025 mol), prepared according to the synthesis procedure described under B1b), in HCl conc. (10 ml) and water (10 ml) was stirred and refluxed for 1 hour. HCl conc. (10 ml) and water (10 ml) were added again. The mixture was stirred and refluxed for 16 hours. The solvent was evaporated. The residue was crystallized from CH<sub>3</sub>OH. The precipitate was filtered off and dried in
- 20 vacuo at 50°C for 16 hours. Yielding: 0.4g of 4-[(4-imidazo[1,2-a]pyridin-3-yl-2-thiazolyl)amino]benzoic acid monohydrochloride (38%) (comp. 7).

#### Example B4

- A mixture of compound 634 (0.0014 mol) in water (60 ml) was stirred and then a
- 25 hydrobromide solution 48% (6 ml) was added. The reaction mixture was stirred and refluxed for 8 hours. The reaction mixture was stirred further for 48 hours at room temperature under N<sub>2</sub> flow. The solvent is evaporated. The residue was crystallized from 2-propanone and CH<sub>3</sub>CN. The precipitate was filtered off and dried. Yield : 0.61g of 6-[2-[[2,3-dichlorophenyl]amino]-4-thiazolyl]pyridinamine monohydrobromide; mp.
- 30 236°C (comp. 635).

#### Example B5


- A mixture of N-[5-[(1-oxo-2-bromo)ethyl]-2-pyridinyl]acetamide (0.002 mol), prepared according to A5c), and [3-(trifluoromethyl)phenyl]thiourea (0.002 mol) in ethanol (100 ml) was stirred and refluxed for 1 hour. The mixture was cooled and the
- 35 precipitate was filtered off. This precipitate was stirred in water (90 ml) and a

hydrobromide solution 48% (10 ml) was added dropwise. The reaction mixture was stirred and refluxed overnight, cooled off and washed with  $\text{CH}_2\text{Cl}_2$  (2x). The aqueous layer was evaporated until dry, stirred in 2-propanone, filtered off and dried. The precipitate was stirred in water and the formed precipitate was filtered off and dried.

- 5 Yield 0.25g of 6-[2-[[3-(trifluoromethyl)phenyl]amino]-4-thiazolyl]pyridinamine monohydrobromide monohydrate; mp. 148°C (comp. 637).

Tables 1 to 12 list the compounds of formula (I) which were prepared according to one of the above described examples.

Table 1



10


Co. no.	Ex. no.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Physical data
8	B1a	CH	H	4-OCH <sub>3</sub>	H	H	HCl (1:1); mp. 235°C
9	B1a	CH	H	H	H	H	HCl (1:1); mp. 170-172°C (dec) *
10	B1a	N	H	4-OCH <sub>3</sub>	H	H	HCl (1:2); mp. 222°C
11	B1a	N	H	H	H	H	HCl (1:2); H <sub>2</sub> O (1:1); mp. 188°C
12	B1a	N	H	3-CF <sub>3</sub>	H	H	HCl (1:2); mp. 190°C
13	B1a	N	H	4-CH <sub>3</sub>	H	H	HCl (1:2); mp. 210°C
14	B1a	N	H	3-CH <sub>3</sub>	H	H	HCl (1:2); mp. 198°C
15	B1a	N	H	3-OCH <sub>3</sub>	H	H	HCl (1:2); mp. 198°C
16	B1a	N	H	4-CF <sub>3</sub>	H	H	HCl (1:1); mp. 228°C
17	B1a	CH	CH <sub>3</sub>	H	H	H	HCl (1:1)
18	B1a	CH	CH <sub>3</sub>	4-OCH <sub>3</sub>	H	H	HCl (1:1)
19	B1a	N	H	4-COOC <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:2)
20	B1a	N	H	4-Br	H	H	HCl (1:2)
22	B1a	CH	H	4-CH <sub>3</sub>	H	H	HCl (1:1); H <sub>2</sub> O (1:1)
48	B1a	N	H	2-F	3-F	4-F	HCl (1:2)
151	B1a	N	H	2-CF <sub>3</sub>	H	H	HCl (1:2)
152	B1a	N	H	4-OCH <sub>2</sub> -phenyl	H	H	HCl (1:2)

Co. no.	Ex. no.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Physical data
153	B1a	N	H	3-Br	H	H	HCl (1:2)
154	B1a	N	H	2-OCF <sub>3</sub>	H	H	HCl (1:2)
155	B1a	N	H	2-CH(CH <sub>3</sub> ) <sub>2</sub>	H	H	HCl (1:2)
156	B1a	N	H	2-SCH <sub>3</sub>	H	H	HCl (1:2)
157	B1a	N	H	2-OC <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:2)
158	B1a	N	H	2-CH <sub>3</sub>	H	H	HCl (1:2)
159	B1a	N	H	2-F	H	H	HCl (1:2)
160	B1a	N	H	3-Cl	4-Br	H	HCl (1:2)
161	B1a	N	H	4-CF <sub>3</sub>	2-Cl	H	HCl (1:2)
162	B1a	N	H	4-CH <sub>3</sub>	3-Cl	H	HCl (1:2)
163	B1a	N	H	2-CH <sub>3</sub>	4-Cl	H	HCl (1:2)
164	B1a	N	H	3-F	4-F	H	HCl (1:2)
165	B1a	N	H	2-CH <sub>3</sub>	3-Cl	H	HCl (1:2)
166	B1a	N	H	2-Cl	3-Cl	H	HCl (1:2); mp. 227-229°C (dec) *
168	B1a	N	H	2-CH <sub>3</sub>	5-Cl	H	HCl (1:2)
169	B1a	N	H	2-CH <sub>3</sub>	5-F	H	HCl (1:2)
170	B1a	N	H	2-CH <sub>3</sub>	4-CH <sub>3</sub>	5-CH <sub>3</sub>	HCl (1:2)
171	B1a	N	H	2-OCH <sub>3</sub>	4-Cl	5-OCH <sub>3</sub>	HCl (1:2)
172	B1a	N	H	2-Cl	4-Cl	5-Cl	HCl (1:2)
173	B1a	N	H	3-OCH <sub>3</sub>	4-OCH <sub>3</sub>	5-OCH <sub>3</sub>	HCl (1:2)
174	B1a	N	H	2-Cl	5-CF <sub>3</sub>	H	HCl (1:2)
175	B1a	N	H	2-OCH <sub>3</sub>	5-Cl	H	HCl (1:2)
176	B1a	N	H	2-OCH <sub>3</sub>	5-CH <sub>3</sub>	H	HCl (1:2)
177	B1a	N	H	2-OCH <sub>3</sub>	5-OCH <sub>3</sub>	H	HCl (1:2)
178	B1a	N	H	3-Cl	5-Cl	H	HCl (1:2)
179	B1a	N	H	2-CH <sub>3</sub>	3-CH <sub>3</sub>	H	HCl (1:2)
180	B1a	N	H	3-CH <sub>3</sub>	5-CH <sub>3</sub>	H	HCl (1:2)
181	B1a	N	H	2-OCH <sub>3</sub>	4-OCH <sub>3</sub>	H	HCl (1:2)
182	B1a	N	H	3-CF <sub>3</sub>	4-Cl	H	HCl (1:2)
183	B1a	N	H	2-Br	4-CH <sub>3</sub>	H	HCl (1:2)
184	B1a	N	H	2-CH <sub>3</sub>	4-CH <sub>3</sub>	H	HCl (1:2)
185	B1a	N	H	2-CF <sub>3</sub>	4-Br	H	HCl (1:2)
187	B1a	N	H	2-OCH <sub>3</sub>	H	H	HCl (1:2)
188	B1a	N	H	2-OH	H	H	HCl (1:2)

Co. no.	Ex. no.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Physical data
189	B1a	N	H	2-Cl	H	H	HCl (1:2)
190	B1a	N	H	2-Br	H	H	HCl (1:2)
191	B1a	N	H	3-SCH <sub>3</sub>	H	H	HCl (1:2)
192	B1a	N	H	3-OH	H	H	HCl (1:2)
193	B1a	N	H	3-F	H	H	HCl (1:2)
194	B1a	N	H	3-CN	H	H	HCl (1:2)
195	B1a	N	H	4-O-phenyl	H	H	HCl (1:2)
196	B1a	N	H	2-(2,4-dichlorophenoxy)	H	H	HCl (1:2)
197	B1a	N	H	2-F	5-F	H	HCl (1:2)
198	B1a	N	H	2-F	4-F	H	HCl (1:2)
199	B1a	N	H	2-Cl	4-Cl	H	HCl (1:2)
200	B1a	N	H	3-Cl	4-Cl	H	HCl (1:2)
203	B1a	N	H	2-C <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:2)
204	B1a	N	H	3-COOH	H	H	HCl (1:2)
205	B1a	N	H	3-COOC <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:2)
206	B1a	N	H	3-COCH <sub>3</sub>	H	H	HCl (1:2)
207	B1a	N	H	4-OH	H	H	HCl (1:2)
208	B1a	N	H	4-OC <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:2)
209	B1a	N	H	4-OCF <sub>3</sub>	H	H	HCl (1:2)
211	B1a	N	H	4-F	H	H	HCl (1:2)
212	B1a	N	H	4-cyclohexyl	H	H	HCl (1:2)
213	B1a	N	H	4-CN	H	H	HCl (1:2)
214	B1a	N	H	4-C <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:2)
215	B1a	N	H	4-COOH	H	H	HCl (1:2)
217	B1a	N	H	3-Cl	H	H	HCl (1:2)
21	B1a	N	H	2-Cl	5-Cl	H	HCl (1:2)
186	B1a	N	H	3-CF <sub>3</sub>	5-CF <sub>3</sub>	H	HCl (1:2)
210	B1a	N	H	3-S(O) <sub>2</sub> -NH <sub>2</sub>	H	H	HCl (1:1)



\* = decomposition

Table 2



Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Physical data
1	B1a	6-Cl	4-OCH <sub>3</sub>	H	H	HCl (1:1)
23	B1a	H	4-OCF <sub>3</sub>	H	H	HCl (1:1); mp. 222°C
24	B1a	H	3-Cl	H	H	HCl (1:1)
25	B1a	H	2-Cl	H	H	HCl (1:1); H <sub>2</sub> O (1:1)
26	B1a	H	3-COOH	H	H	HCl (1:1)
27	B1a	H	3-COOC <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:1)
28	B1a	H	2-OCH <sub>3</sub>	4-OCH <sub>3</sub>	H	HCl (1:1); mp. 158°C
29	B1a	H	3-OCH <sub>3</sub>	H	H	HCl (1:1)
30	B1a	H	3-Cl	5-Cl	H	HCl (1:1)
31	B1a	H	3-CH <sub>3</sub>	H	H	HCl (1:1); mp. 218-220°C (dec) *
32	B1a	H	4-OC <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:1)
33	B1a	H	3-S-CH <sub>3</sub>	H	H	HCl (1:1); mp. 220°C
34	B1a	H	2-OCH <sub>3</sub>	H	H	HCl (1:1); ethanolate (1:1); mp. 152°C
35	B1a	H	3-OH	H	H	HCl (1:1)
36	B1a	H	3-COCH <sub>3</sub>	H	H	HCl (1:1)
37	B1a	H	4-Cl	H	H	HCl (1:1); ethanolate (1:1)
38	B1a	H	3-CF <sub>3</sub>	4-Cl	H	HCl (1:1)
39	B1a	H	4-CH <sub>3</sub>	H	H	HCl (1:1); mp. >250°C
40	B1a	H	2-OH	H	H	HCl (1:2)
41	B1a	H	2-S-CH <sub>3</sub>	H	H	HCl (1:2); ethanolate (1:1)
42	B1a	H	4-I	H	H	HCl (1:1)
43	B1a	H	3-Cl	4-Cl	H	HCl (1:1); mp. >260°C
44	B1a	H	4-COOC <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:1)
45	B1a	H	2-Cl	3-Cl	H	HCl (1:1), H <sub>2</sub> O (1:1); mp. 150-154°C (dec)*
46	B1a	H	2-F	3-F	4-F	HCl (1:1)

Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Physical data
47	B1a	H	3-CH <sub>3</sub>	5-CH <sub>3</sub>	H	HCl (1:1)
52	B1a	H	4-cyclohexyl	H	H	HCl (1:1)
54	B1a	6-Cl	4-CH <sub>3</sub>	H	H	HCl (1:1); mp. 232°C
55	B1a	6-Cl	3-CF <sub>3</sub>	H	H	HCl (1:1); H <sub>2</sub> O (1:1); mp. 222°C
56	B1a	6-Cl	3-OH	H	H	HCl (1:1)
57	B1a	H	4-CH(CH <sub>3</sub> ) <sub>2</sub>	H	H	HCl (1:1)
58	B1a	H	2-Cl	4-Cl	6-Cl	HCl (1:1)
59	B1a	H	2-Cl	6-Cl	H	
60	B1a	H	2-CH <sub>3</sub>	6-CH <sub>3</sub>	H	
61	B1b	2-CH <sub>3</sub>	4-OCH <sub>3</sub>	H	H	
62	B1b	2-CH <sub>3</sub>	H	H	H	mp. 221-223°C
63	B1b	2-CH <sub>3</sub>	2-CH <sub>3</sub>	H	H	HBr (1:1); mp. 190°C
64	B1b	2-CH <sub>3</sub>	4-CH <sub>3</sub>	H	H	HBr (1:1); ethanolate (1:1); mp. >260°C
65	B1b	2-CH <sub>3</sub>	2-F	H	H	HBr (1:1); mp. 246°C
66	B1b	2-CH <sub>3</sub>	3-F	H	H	HBr (1:1)
67	B1b	2-CH <sub>3</sub>	4-F	H	H	HBr (1:1); mp. >258°C
68	B1b	2-CH <sub>3</sub>	3-CN	H	H	HBr (1:1)
69	B1b	2-CH <sub>3</sub>	4-CN	H	H	HBr (1:1); mp. >260°C
70	B1b	2-CH <sub>3</sub>	2-OCH <sub>3</sub>	H	H	HBr (1:1); mp. 204°C
71	B1b	2-CH <sub>3</sub>	4-OH	H	H	HBr (1:1); mp. >260°C
72	B1b	2-CH <sub>3</sub>	2-CF <sub>3</sub>	H	H	HBr (1:1); mp. 250°C
73	B1b	2-CH <sub>3</sub>	2-OCF <sub>3</sub>	H	H	HBr (1:1)
74	B1b	H	4-OCH <sub>3</sub>	H	H	HBr (1:1); mp. 254°C
75	B1b	H	4-CH <sub>3</sub>	H	H	HBr (1:1); mp. >260°C
76	B1b	2-CH <sub>3</sub>	3-CF <sub>3</sub>	H	H	HBr (1:1); mp. 256°C
77	B1b	2-CH <sub>3</sub>	4-OCF <sub>3</sub>	H	H	HBr (1:1)
78	B1b	2-CH <sub>3</sub>	4-CF <sub>3</sub>	H	H	HBr (1:1); ethanolate (1:1)
79	B1b	H	2-F	H	H	HBr (1:1)
80	B1b	H	H	H	H	HBr (1:1)
81	B1b	H	3-CF <sub>3</sub>	H	H	HBr (1:1); mp. 260- 262°C (dec) *


Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Physical data
82	B1b	H	4-CF <sub>3</sub>	H	H	HBr (1:1); mp. 260-262°C (dec) *
83	B1b	H	4-OH	H	H	HBr (1:1); mp. >260°C
84	B1b	2-CH <sub>3</sub>	4-S-CH <sub>3</sub>	H	H	HBr (1:1)
85	B1b	H	4-S-CH <sub>3</sub>	H	H	HBr (1:1)
86	B1b	H	4-C <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1); mp. >260°C
87	B1b	2-CH <sub>3</sub>	4-C <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1); mp. 238°C
90	B1b	H	4-F	H	H	HCl (1:1)
92	B1b	H	3-CN	H	H	
6	B1b	H	4-CN	H	H	HBr (1:1)
99	B1b	2-CH <sub>3</sub>	2-Cl	H	H	HBr (1:1)
104	B1b	2-CH <sub>3</sub>	3-Cl	H	H	HBr (1:1)
105	B1b	H	3-CF <sub>3</sub>	5-CF <sub>3</sub>	H	
107	B1b	H	2-OCF <sub>3</sub>	H	H	HBr (1:1); mp. >250°C
124	B1b	7-Cl	4-OCH <sub>3</sub>	H	H	HBr (1:1)
5	B2	H	3-NH <sub>2</sub>	H	H	HCl (1:2); H <sub>2</sub> O (1:1)
7	B3	H	4-COOH	H	H	HCl (1:1)
134	B1b	2-CH <sub>3</sub>	4-OCH <sub>3</sub>	H	H	HBr (1:2)
135	B1b	2-CH <sub>3</sub>	4-Br	H	H	HBr (1:1)
136	B1b	2-CH <sub>3</sub>	3-CH <sub>3</sub>	H	H	HBr (1:1)
137	B1c	2-CH <sub>3</sub>	4-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	HBr (1:1)
138	B1b	2-CH <sub>3</sub>	4- 	H	H	HBr (1:1); H <sub>2</sub> O(1:1)
139	B1c	2-CH <sub>3</sub>	4-NH <sub>2</sub>	H	H	HBr (1:1)
141	B1b	2-CH <sub>3</sub>	4-NH-CO-CH <sub>3</sub>	H	H	HBr (1:1)
142	B1b	2-CH <sub>3</sub>	3-OH	H	H	HBr (1:1)
144	B1b	2-CH <sub>3</sub>	4- 	H	H	HBr (1:1)
465	B1a	H	4-OCH <sub>2</sub> -phenyl	H	H	HCl (1:1)
466	B1a	H	3-Br	H	H	

Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Physical data
467	B1a	H	2-OCF <sub>3</sub>	H	H	HCl (1:1)
468	B1a	H	2-CH(CH <sub>3</sub> ) <sub>2</sub>	H	H	HCl (1:1)
469	B1a	H	2-SCH <sub>3</sub>	H	H	HCl (1:1)
470	B1a	H	2-OC <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:1)
471	B1a	H	2-CH <sub>3</sub>	H	H	HCl (1:1)
472	B1a	H	2-F	H	H	HCl (1:1)
473	B1a	H	2-CF <sub>3</sub>	H	H	HCl (1:1)
474	B1a	H	3-Cl	4-Br	H	HCl (1:1)
475	B1a	H	2-Cl	4-CF <sub>3</sub>	H	HCl (1:1)
476	B1a	H	3-Cl	4-CH <sub>3</sub>	H	HCl (1:1)
477	B1a	H	2-CH <sub>3</sub>	4-Cl	H	HCl (1:1)
478	B1a	H	3-F	4-F	H	HCl (1:1)
479	B1a	H	2-CH <sub>3</sub>	3-Cl	H	HCl (1:1)
481	B1a	H	2-CH <sub>3</sub>	5-Cl	H	HCl (1:1)
482	B1a	H	2-CH <sub>3</sub>	5-F	H	HCl (1:1)
483	B1a	H	2-CH <sub>3</sub>	4-CH <sub>3</sub>	5-CH <sub>3</sub>	HCl (1:1)
484	B1a	H	2-OCH <sub>3</sub>	4-Cl	5-OCH <sub>3</sub>	HCl (1:1)
485	B1a	H	2-Cl	4-Cl	5-Cl	HCl (1:1)
486	B1a	H	3-OCH <sub>3</sub>	4-OCH <sub>3</sub>	5-OCH <sub>3</sub>	HCl (1:1)
488	B1a	H	2-Cl	5-CF <sub>3</sub>	H	HCl (1:1)
489	B1a	H	2-OCH <sub>3</sub>	5-Cl	H	HCl (1:1)
490	B1a	H	2-OCH <sub>3</sub>	5-CH <sub>3</sub>	H	HCl (1:1)
491	B1a	H	2-OCH <sub>3</sub>	5-OCH <sub>3</sub>	H	HCl (1:1)
492	B1a	H	2-CH <sub>3</sub>	3-CH <sub>3</sub>	H	HCl (1:1)
494	B1a	H	2-Br	4-CH <sub>3</sub>	H	HCl (1:1)
495	B1a	H	2-CH <sub>3</sub>	4-CH <sub>3</sub>	H	HCl (1:1)
496	B1a	H	2-CF <sub>3</sub>	4-Br	H	HCl (1:1)
498	B1a	H	2-Br	H	H	HCl (1:1)
499	B1a	H	3-F	H	H	HCl (1:1)
500	B1a	H	3-CN	H	H	HCl (1:1)
501	B1a	H	4-phenoxy	H	H	HCl (1:1)
502	B1a	H	2-C <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:1)
504	B1a	H	2-Cl	4-Cl	H	HCl (1:1)
505	B1a	H	2-F	4-F	H	HCl (1:1)
506	B1a	H	2-Cl	5-Cl	H	HCl (1:1)







Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Physical data
507	B1a	H	2-F	5-F	H	HCl (1:1)
508	B1a	H	2-(2,4-dichlorophenoxy)	H	H	HCl (1:1)
509	B1a	H	4-Br	H	H	HCl (1:1)
4	B1b	7-Cl	3-CF <sub>3</sub>	H	H	HBr (1:1)
480	B1a	H	4-S(O) <sub>2</sub> -NH <sub>2</sub>	H	H	HCl (1:1)
497	B1a	H	3-S(O) <sub>2</sub> -NH <sub>2</sub>	H	H	HCl (1:1)
223	B1a	H	3-S(O) <sub>2</sub> -CH <sub>3</sub>	H	H	HCl (1:1)
239	B1a	H	3-CH <sub>2</sub> -OH	H	H	HCl (1:1)
244	B1a	H	3-O-CH <sub>3</sub>	4-O-CH <sub>3</sub>	H	HCl (1:1)
254	B1a	H	3-CF <sub>3</sub>	H	H	HCl (1:1)
265	B1a	H	4-CF <sub>3</sub>	H	H	HCl (1:1)
291	B1a	H	4-N <sub>3</sub>	H	H	
299	B1a	H	4-C(=O)-CH <sub>3</sub>	H	H	HCl (1:1)
311	B1a	H	3-CH <sub>3</sub>	4-F	H	HCl (1:1); mp. 250-252°C (dec) *



\* = decomposition

Table 3




Co. no.	Ex. no.	R <sup>1</sup>	Het	Physical data
88	B1b	H	4-pyridinyl	HBr (1:1)
89	B1b	H	2-thiazolyl	HBr (1:1)
91	B1b	H	1H-pyrazol-3-yl	HBr (1:1); mp. 188°C
93	B1b	H	3-benzo[b]furanyl	HBr (1:1)
94	B1b	4-OCH <sub>3</sub>	3-benzo[b]furanyl	HBr (1:1)
96	B1b	4-OCH <sub>3</sub>	4-pyridinyl	HBr (1:1); ethanolate (1:1); mp. 250°C
97	B1b	4-OCH <sub>3</sub>	1H-pyrazol-3-yl	HBr (1:1)
98	B1b	4-OCH <sub>3</sub>	2-thiazolyl	HBr (1:2)
100	B1b	3-CF <sub>3</sub>	3-quinolinyl	HBr (1:1); H <sub>2</sub> O (1:1); mp. 171-173°C (dec) *











Co. no.	Ex. no.	R <sup>1</sup>	Het	Physical data
102	B1b	3-CF <sub>3</sub>	4-pyridinyl	HBr (1:1); mp. >250°C
103	B1b	3-CF <sub>3</sub>	2-thiazolyl	HBr (1:1); mp. 222°C
106	B1b	4-OCH <sub>3</sub>	3-quinolinyl	HBr (1:1); H <sub>2</sub> O (1:1)
108	B1b	4-OCH <sub>3</sub>	1 <i>H</i> -indazol-3-yl	HBr (1:1); mp. 212°C
111	B1b	3-CF <sub>3</sub>	1 <i>H</i> -indazol-3-yl	HBr (1:1)
112	B1b	H	1 <i>H</i> -indazol-3-yl	HBr (1:1); mp. 238°C
113	B1b	4-CH <sub>3</sub>		
119	B1b	3-CH <sub>3</sub>		HBr (1:1); mp. 202-204°C (dec) *
120	B1b	4-Br		HBr (1:1)
125	B1c	4-OCH <sub>3</sub>		HBr (1:1)
140	B1b	4-CH <sub>3</sub>		HBr (1:2); mp. >260°C
145	B1b	4-OC <sub>2</sub> H <sub>5</sub>	2,4-dimethyl-5-thiazolyl	
146	B1b	4-SO <sub>2</sub> -NH <sub>2</sub>	2-amino-4-methyl-5-thiazolyl	
332	B1c	3-CF <sub>3</sub>		HBr (1:1)
359	B1a	H	4-pyridinyl	
373	B1a	3-CH <sub>3</sub>	4-pyridinyl	
387	B1a	4-NH <sub>2</sub>	4-pyridinyl	HBr (1:1)
427	B1a	4-CH <sub>3</sub>	4-pyridinyl	
437	B1a	4-O-C <sub>2</sub> H <sub>5</sub>	4-pyridinyl	
449	B1a	3-OH	4-pyridinyl	
511	B1b	3-CF <sub>3</sub>	1,2-dimethyl-1 <i>H</i> -imidazol-5-yl	HBr (1:1)
512	B1b	3-Cl	4-pyridinyl	HBr (1:1)
513	B1a	H	5-chloro-2-thienyl	
514	B1a	4-Br	2,4-dimethyl-	

Co. no.	Ex. no.	R <sup>1</sup>	Het	Physical data
			5-thiazolyl	
515	B1a	3-CH <sub>3</sub>		mp 214°C HCl (1:1); mp. 120-122°C (dec) * HCl (1:1)
516	B1a	3-OH		
517	B1b	3-CF <sub>3</sub>	5-pyrimidinyl	
518	B1b	3-CF <sub>3</sub>	3-furanyl	
519	B1b	3-CF <sub>3</sub>	2-furanyl	

\* = decomposition

Table 4



Co. no.	Ex. no.	Het	Q	Physical data
130	B1a			HCl (1:1)
131	B1b			HBr (1:1)
132	B1b		3-pyridinyl	HBr (1:1); H <sub>2</sub> O (1:1) HBr (1:1)
133	B1c		3-pyridinyl	
143	B1c		2-pyridinyl	
150	B1b		cyclohexyl	
201	B1a			












Co. no.	Ex. no.	Het	Q	Physical data
202	B1a		1-naphthalenyl	HCl (1:2)
237	B1b			HBr (1:1)
238	B1b		1-naphthalenyl	HBr (1:1)
426	B1a		1-naphthalenyl	HCl (1:1)
510	B1a		1-naphthalenyl	HBr (1:1)
218	B1a		cyclohexyl	HCl (1:1)
520	B1a	4-pyridinyl	2,6-dichlorophenyl	
521	B1a	4-pyridinyl	2,6-dimethylphenyl	
522	B1a		3-pyridinyl	HBr (1:2)
523	B1a	5-chloro-2-thienyl	3-pyridinyl	HBr (1:1)
524	B1a		2-pyridinyl	
525	B1b	3-furanyl	2,3-dichlorophenyl	
526	B1a		6-methoxy-3-pyridinyl	mp. 210-212°C (dec) *
527	B1a	1,2-dimethyl-1H-imidazol-5-yl	6-chloro-3-pyridinyl	HBr (1:1); mp. 174-176°C (dec)


Table 5



Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Physical data
2	B1b	4-Cl	H	H	HBr (1:1)
114	B1b	4-OCH <sub>3</sub>	H	H	HBr (1:1)
115	B1b	3-CH <sub>3</sub>	H	H	HBr (1:1); H <sub>2</sub> O (1:1)
116	B1b	4-CH <sub>3</sub>	H	H	HBr (1:1); H <sub>2</sub> O (1:1)
117	B1b	H	H	H	HBr (1:1); H <sub>2</sub> O (1:1)
118	B1b	3-OH	H	H	HBr (1:1)
121	B1b	3-CF <sub>3</sub>	H	H	
123	B1b	4-Br	H	H	HBr (1:1)
219	B1b	2-Cl	4-Cl	5-Cl	HBr (1:1)
220	B1b	2-OCH <sub>3</sub>	4-Cl	5-OCH <sub>3</sub>	HBr (1:1)
221	B1b	2-CH <sub>3</sub>	4-CH <sub>3</sub>	5-CH <sub>3</sub>	HBr (1:1)
222	B1b	2-CH <sub>3</sub>	5-Cl	H	HBr (1:1)
224	B1b	2-Cl	5-Cl	H	HBr (1:1)
225	B1b	2-OCH <sub>3</sub>	5-Cl	H	HBr (1:1)
226	B1b	2-OCH <sub>3</sub>	5-CH <sub>3</sub>	H	HBr (1:1)
227	B1b	2-OCH <sub>3</sub>	5-OCH <sub>3</sub>	H	HBr (1:1)
228	B1b	3-Cl	5-Cl	H	HBr (1:1)
229	B1b	3-CH <sub>3</sub>	5-CH <sub>3</sub>	H	HBr (1:1)
230	B1b	2-OCH <sub>3</sub>	4-OCH <sub>3</sub>	H	HBr (1:1)
231	B1b	3-F	4-F	H	HBr (1:1)
232	B1b	2-Cl	4-Cl	H	HBr (1:1)
233	B1b	2-CH <sub>3</sub>	4-Cl	H	HBr (1:1)
234	B1b	3-Cl	4-Cl	H	HBr (1:1)
235	B1b	3-CF <sub>3</sub>	4-Cl	H	HBr (1:1)
236	B1b	4-CH <sub>3</sub>	3-Cl	H	HBr (1:1)
240	B1b	2-OCH <sub>3</sub>	H	H	HBr (1:1)
241	B1b	2-OH	H	H	HBr (1:1)
242	B1b	2-Br	H	H	HBr (1:1)
243	B1b	3-SCH <sub>3</sub>	H	H	HBr (1:1)

Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Physical data
245	B1b	3-F	H	H	HBr (1:1)
246	B1b	3-CN	H	H	HBr (1:1)
247	B1b	3-Cl	H	H	HBr (1:1)
248	B1b	3-COOH	H	H	HBr (1:1)
249	B1b	3-COOC <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
250	B1b	3-Br	H	H	HBr (1:1)
251	B1b	4-OH	H	H	HBr (1:1)
252	B1b	4-phenoxy	H	H	HBr (1:1)
253	B1b	4-OCH <sub>2</sub> -phenyl	H	H	HBr (1:1)
255	B1b	4-F	H	H	HBr (1:1)
256	B1b	4-cyclohexyl	H	H	HBr (1:1)
257	B1b	4-COOH	H	H	HBr (1:1)
258	B1b	4-COOC <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
259	B1b	4-COCH <sub>3</sub>	H	H	HBr (1:1)
260	B1b	4-OC <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
261	B1b	4-C <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)

Table 6




Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Physical data
262	B1b	2-(2,4-dichlorophenoxy)	H	H	HBr (1:1)
263	B1b	3-OCH <sub>3</sub>	4-OCH <sub>3</sub>	5-OCH <sub>3</sub>	HBr (1:1)
264	B1b	2-CH <sub>3</sub>	4-CH <sub>3</sub>	5-CH <sub>3</sub>	HBr (1:1)
266	B1b	2-F	5-F	H	HBr (1:1)
267	B1b	2-Cl	5-Cl	H	HBr (1:1)
268	B1b	2-Cl	5-CF <sub>3</sub>	H	HBr (1:1)
269	B1b	2-OCH <sub>3</sub>	5-Cl	H	HBr (1:1)
270	B1b	2-OCH <sub>3</sub>	5-CH <sub>3</sub>	H	HBr (1:1)
271	B1b	2-OCH <sub>3</sub>	5-OCH <sub>3</sub>	H	HBr (1:1)
272	B1b	3-Cl	5-Cl	H	HBr (1:1)

Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Physical data
273	B1b	2-Cl	3-Cl	H	HBr (1:1)
274	B1b	2-CH <sub>3</sub>	3-Cl	H	HBr (1:1)
275	B1b	3-CH <sub>3</sub>	5-CH <sub>3</sub>	H	HBr (1:1)
276	B1b	3-CF <sub>3</sub>	5-CF <sub>3</sub>	H	HBr (1:1)
277	B1b	2-OCH <sub>3</sub>	4-OCH <sub>3</sub>	H	HBr (1:1)
278	B1b	2-F	4-F	H	HBr (1:1)
279	B1b	3-F	4-F	H	HBr (1:1)
280	B1b	2-Cl	4-Cl	H	HBr (1:1)
281	B1b	2-CH <sub>3</sub>	4-Cl	H	HBr (1:1)
282	B1b	3-Cl	4-Cl	H	HBr (1:1)
283	B1b	3-CF <sub>3</sub>	4-Cl	H	HBr (1:1)
284	B1b	2-Br	4-CH <sub>3</sub>	H	HBr (1:1)
285	B1b	2-CH <sub>3</sub>	4-CH <sub>3</sub>	H	HBr (1:1)
286	B1b	3-Cl	4-CH <sub>3</sub>	H	HBr (1:1)
287	B1b	2-Cl	4-CF <sub>3</sub>	H	HBr (1:1)
288	B1b	2-CF <sub>3</sub>	4-Br	H	HBr (1:1)
289	B1b	3-Cl	4-Br	H	HBr (1:1)
292	B1b	2-OCH <sub>3</sub>	H	H	HBr (1:1)
293	B1b	2-OH	H	H	HBr (1:1)
294	B1b	2-Cl	H	H	HBr (1:1)
295	B1b	2-F	H	H	HBr (1:1)
296	B1b	2-CF <sub>3</sub>	H	H	HBr (1:1)
297	B1b	3-SCH <sub>3</sub>	H	H	HBr (1:1)
298	B1b	3-OH	H	H	HBr (1:1)
300	B1b	3-F	H	H	HBr (1:1)
301	B1b	3-CN	H	H	HBr (1:1)
302	B1b	3-Cl	H	H	HBr (1:1)
303	B1b	3-COOH	H	H	HBr (1:1)
304	B1b	3-COOC <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
305	B1b	3-COCH <sub>3</sub>	H	H	HBr (1:1)
306	B1b	3-Br	H	H	HBr (1:1)
307	B1b	4-phenoxy	H	H	HBr (1:1)
308	B1b	4-OC <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
309	B1b	4-OCF <sub>3</sub>	H	H	HBr (1:1)
310	B1b	4-OCH <sub>2</sub> -phenyl	H	H	HBr (1:1)

Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Physical data
312	B1b	4-F	H	H	HBr (1:1)
313	B1b	4-cyclohexyl	H	H	HBr (1:1)
314	B1b	4-Cl	H	H	HBr (1:1)
315	B1b	4-C <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
316	B1b	4-COOH	H	H	HBr (1:1)
317	B1b	4-COOC <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
319	B1b	2-SCH <sub>3</sub>	H	H	HBr (1:1)
320	B1b	2-OCF <sub>3</sub>	H	H	HBr (1:1)
321	B1b	2-Br	H	H	HBr (1:1)
322	B1b	2-C <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
323	B1b	2-CH <sub>3</sub>	3-CH <sub>3</sub>	H	HBr (1:1)
528	B1b	2-F	3-F	4-F	HBr (1:1)
529	B1b	2-Cl	4-Cl	5-Cl	HBr (1:1)
530	B1b	2-OCH <sub>3</sub>	4-Cl	5-OCH <sub>3</sub>	HBr (1:1)
531	B1b	2-CH <sub>3</sub>	5-F	H	HBr (1:1)
532	B1b	2-CH <sub>3</sub>	5-Cl	H	HBr (1:1)
533	B1b	2-O-C <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
534	B1c	3-CH <sub>3</sub>	H	H	HCl (1:1)
535	B1b	4-CF <sub>3</sub>	H	H	HBr (1:1)
536	B1b	3-CF <sub>3</sub>	H	H	HBr (1:1)

Table 7



Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Physical data
95	B1b	4-OCH <sub>3</sub>	H	H	HBr (1:2); mp. 228°C
101	B1b	3-CF <sub>3</sub>	H	H	HBr (1:1); mp. 238°C
122	B1b	4-Br	H	H	HBr (1:1)
147	B1b	4-OCH <sub>3</sub>	H	H	
148	B1b	3-OH	H	H	
149	B1b	4-SO <sub>2</sub> -NH <sub>2</sub>	H	H	HBr (1:1)


Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Physical data
324	B1b	2-(2,4-dichlorophenoxy)	H	H	HBr (1:1)
325	B1b	3-OCH <sub>3</sub>	4-OCH <sub>3</sub>	5-OCH <sub>3</sub>	HBr (1:1)
326	B1b	2-F	3-F	4-F	HBr (1:1)
327	B1b	2-Cl	4-Cl	5-Cl	HBr (1:1)
328	B1b	2-OCH <sub>3</sub>	4-Cl	5-OCH <sub>3</sub>	HBr (1:1)
329	B1b	2-CH <sub>3</sub>	4-CH <sub>3</sub>	5-CH <sub>3</sub>	HBr (1:1)
330	B1b	2-CH <sub>3</sub>	5-F	H	HBr (1:1)
331	B1b	2-CH <sub>3</sub>	5-Cl	H	HBr (1:1)
333	B1b	2-F	5-F	H	HBr (1:1)
334	B1b	2-Cl	5-Cl	H	HBr (1:1)
335	B1b	2-OCH <sub>3</sub>	5-Cl	H	HBr (1:1)
336	B1b	2-OCH <sub>3</sub>	5-CH <sub>3</sub>	H	HBr (1:1)
337	B1b	2-OCH <sub>3</sub>	5-OCH <sub>3</sub>	H	HBr (1:1)
338	B1b	3-Cl	5-Cl	H	HBr (1:1)
339	B1b	2-Cl	3-Cl	H	HBr (1:1)
340	B1b	2-CH <sub>3</sub>	3-Cl	H	HBr (1:1)
341	B1b	2-CH <sub>3</sub>	3-CH <sub>3</sub>	H	HBr (1:1)
342	B1b	3-CH <sub>3</sub>	5-CH <sub>3</sub>	H	HBr (1:1)
343	B1b	3-CF <sub>3</sub>	5-CF <sub>3</sub>	H	HBr (1:1)
344	B1b	2-OCH <sub>3</sub>	4-OCH <sub>3</sub>	H	HBr (1:1)
345	B1b	2-F	4-F	H	HBr (1:1)
346	B1b	3-F	4-F	H	HBr (1:1)
347	B1b	2-Cl	4-Cl	H	HBr (1:1)
348	B1b	2-CH <sub>3</sub>	4-Cl	H	HBr (1:1)
349	B1b	3-Cl	4-Cl	H	HBr (1:1)
350	B1b	3-CF <sub>3</sub>	4-Cl	H	HBr (1:1)
351	B1b	2-Br	4-CH <sub>3</sub>	H	HBr (1:1)
352	B1b	2-CH <sub>3</sub>	4-CH <sub>3</sub>	H	HBr (1:1)
353	B1b	3-Cl	4-CH <sub>3</sub>	H	HBr (1:1)
354	B1b	2-Cl	4-CF <sub>3</sub>	H	HBr (1:1)
355	B1b	2-CF <sub>3</sub>	4-Br	H	HBr (1:1)
356	B1b	3-Cl	4-Br	H	HBr (1:1)
360	B1b	2-OCH <sub>3</sub>	H	H	HBr (1:1)
361	B1b	2-OH	H	H	HBr (1:1)
362	B1b	2-Cl	H	H	HBr (1:1)

Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Physical data
363	B1b	2-F	H	H	HBr (1:1)
364	B1b	2-CF <sub>3</sub>	H	H	HBr (1:1)
365	B1b	2-C <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
366	B1b	2-OC <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
367	B1b	2-SCH <sub>3</sub>	H	H	HBr (1:1)
368	B1b	2-CH(CH <sub>3</sub> ) <sub>2</sub>	H	H	HBr (1:1)
369	B1b	2-OCF <sub>3</sub>	H	H	HBr (1:1)
370	B1b	2-Br	H	H	HBr (1:1)
371	B1b	3-SCH <sub>3</sub>	H	H	HBr (1:1)
372	B1b	3-OCH <sub>3</sub>	H	H	HBr (1:1)
374	B1b	3-F	H	H	HBr (1:1)
375	B1b	3-CN	H	H	HBr (1:1)
376	B1b	3-Cl	H	H	HBr (1:1)
377	B1b	3-CH <sub>3</sub>	H	H	HBr (1:1)
378	B1b	3-COOH	H	H	HBr (1:1)
379	B1b	3-COOC <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
380	B1b	3-COCH <sub>3</sub>	H	H	HBr (1:1)
381	B1b	3-Br	H	H	HBr (1:1)
382	B1b	4-OH	H	H	HBr (1:1)
383	B1b	4-phenoxy	H	H	HBr (1:1)
384	B1b	4-OC <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
385	B1b	4-OCF <sub>3</sub>	H	H	HBr (1:1)
386	B1b	4-OCH <sub>2</sub> -phenyl	H	H	HBr (1:1)
388	B1b	4-F	H	H	HBr (1:1)
389	B1b	4-cyclohexyl	H	H	HBr (1:1)
390	B1b	4-Cl	H	H	HBr (1:1)
391	B1b	4-C <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
392	B1b	4-CF <sub>3</sub>	H	H	HBr (1:1)
393	B1b	4-COOH	H	H	HBr (1:1)
394	B1b	4-COOC <sub>2</sub> H <sub>5</sub>	H	H	HBr (1:1)
395	B1b	4-COCH <sub>3</sub>	H	H	HBr (1:1)
537	B1a	H	H	H	
538	B1b	3-[SO <sub>2</sub> -NH <sub>2</sub> ]	H	H	HBr (1:1)
539	B1b	3-CH <sub>2</sub> -OH	H	H	HBr (1:1)
540	B1b	3-OCH <sub>3</sub>	4-OCH <sub>3</sub>		HBr (1:2)

Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Physical data
541	B1b	4-CH <sub>3</sub>	H	H	HBr (1:2)
542	B1b	2-Cl	3-Cl	H	
543	B1b	3-CH <sub>3</sub>	4-F	H	HBr (1:1); mp. 242-244°C (dec) *

\* = decomposition

Table 8




Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Physical data
216	B1a	4-CH <sub>3</sub>	H	H	HCl (1:1)
49	B1a	3-CF <sub>3</sub>	H	H	HCl (1:1); mp. 246°C
50	B1a	H	H	H	HCl (1:1); mp. 228°C
51	B1a	3-OCH <sub>3</sub>	H	H	HCl (1:1); mp. 214°C
53	B1a	4-Br	H	H	HCl (1:1)
396	B1a	2-(2,4-dichlorophenoxy)	H	H	HCl (1:1)
397	B1a	2-F	3-F	4-F	HCl (1:1)
398	B1a	2-Cl	4-Cl	5-Cl	HCl (1:1)
399	B1a	2-OCH <sub>3</sub>	4-Cl	5-OCH <sub>3</sub>	HCl (1:1)
400	B1a	2-CH <sub>3</sub>	4-CH <sub>3</sub>	5-CH <sub>3</sub>	HCl (1:1)
401	B1a	2-CH <sub>3</sub>	5-F	H	HCl (1:1)
402	B1a	2-CH <sub>3</sub>	5-Cl	H	HCl (1:1)
404	B1a	2-F	5-F	H	HCl (1:1)
405	B1a	2-Cl	5-Cl	H	HCl (1:1)
406	B1a	2-Cl	5-CF <sub>3</sub>	H	HCl (1:1)
407	B1a	2-OCH <sub>3</sub>	5-Cl	H	HCl (1:1)
408	B1a	2-OCH <sub>3</sub>	5-OCH <sub>3</sub>	H	HCl (1:1)
409	B1a	3-Cl	5-Cl	H	HCl (1:1)
410	B1a	2-Cl	3-Cl	H	HCl (1:1); mp. 202-204°C (dec) *
411	B1a	2-CH <sub>3</sub>	3-Cl	H	HCl (1:1)

Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Physical data
412	B1a	3-CH <sub>3</sub>	5-CH <sub>3</sub>	H	HCl (1:1)
413	B1a	3-CF <sub>3</sub>	5-CF <sub>3</sub>	H	HCl (1:1)
414	B1a	2-F	4-F	H	HCl (1:1)
415	B1a	3-F	4-F	H	HCl (1:1)
416	B1a	2-Cl	4-Cl	H	HCl (1:1)
417	B1a	2-CH <sub>3</sub>	4-Cl	H	HCl (1:1)
418	B1a	3-Cl	4-Cl	H	HCl (1:1)
419	B1a	3-CF <sub>3</sub>	4-Cl	H	HCl (1:1)
420	B1a	2-Br	4-CH <sub>3</sub>	H	HCl (1:1)
421	B1a	2-CH <sub>3</sub>	4-CH <sub>3</sub>	H	HCl (1:1)
422	B1a	3-Cl	4-CH <sub>3</sub>	H	HCl (1:1)
423	B1a	2-Cl	4-CF <sub>3</sub>	H	HCl (1:1)
424	B1a	2-CF <sub>3</sub>	4-Br	H	HCl (1:1)
425	B1a	3-Cl	4-Br	H	HCl (1:1)
428	B1a	2-OCH <sub>3</sub>	H	H	HCl (1:1)
429	B1a	2-OH	H	H	HCl (1:1)
430	B1a	2-Cl	H	H	HCl (1:1)
431	B1a	2-F	H	H	HCl (1:1)
432	B1a	2-CH <sub>3</sub>	H	H	HCl (1:1)
433	B1a	2-OC <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:1)
434	B1a	2-SCH <sub>3</sub>	H	H	HCl (1:1)
435	B1a	2-OCF <sub>3</sub>	H	H	HCl (1:1)
436	B1a	3-SCH <sub>3</sub>	H	H	HCl (1:1)
438	B1a	3-F	H	H	HCl (1:1)
439	B1a	3-CN	H	H	HCl (1:1)
440	B1a	3-Cl	H	H	HCl (1:1)
441	B1a	3-COOH	H	H	HCl (1:1)
442	B1a	3-COOC <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:1)
443	B1a	3-COCH <sub>3</sub>	H	H	HCl (1:1)
444	B1a	3-Br	H	H	HCl (1:1)
445	B1a	4-OH	H	H	HCl (1:1)
446	B1a	4-OC <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:1)
447	B1a	4-OCF <sub>3</sub>	H	H	HCl (1:1)
448	B1a	4-OCH <sub>2</sub> -phenyl	H	H	HCl (1:1)
450	B1a	4-F	H	H	HCl (1:1)

Co. no.	Ex. no.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	Physical data
451	B1a	4-cyclohexyl	H	H	HCl (1:1)
452	B1a	4-C <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:1)
453	B1a	4-COOH	H	H	HCl (1:1)
454	B1a	4-COOC <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:1)
455	B1a	4-COCH <sub>3</sub>	H	H	HCl (1:1)
456	B1a	3-OCH <sub>3</sub>	4-OCH <sub>3</sub>	5-OCH <sub>3</sub>	HCl (1:1)
457	B1a	2-OCH <sub>3</sub>	5-CH <sub>3</sub>	H	HCl (1:1)
458	B1a	2-CH <sub>3</sub>	3-CH <sub>3</sub>	H	HCl (1:1)
459	B1a	2-OCH <sub>3</sub>	4-OCH <sub>3</sub>	H	HCl (1:1)
460	B1a	2-CF <sub>3</sub>	H	H	HCl (1:1)
461	B1a	2-C <sub>2</sub> H <sub>5</sub>	H	H	HCl (1:1)
462	B1a	2-CH(CH <sub>3</sub> ) <sub>2</sub>	H	H	HCl (1:1)
463	B1a	2-Br	H	H	HCl (1:1)
464	B1a	4-phenoxy	H	H	HCl (1:1)
544	B1a	4-OCH <sub>3</sub>	H	H	HCl (1:1)
545	B1a	3-OH	H	H	HCl (1:1); H <sub>2</sub> O (1:1); mp. 186°C

\* = decomposition

Table 9




5

Co. no.	Ex. no.	Q	Physical data
546	B1a	6-chloro-3-pyridinyl	HCl (1:1)
547	B1a	3-pyridinyl	HCl (1:2); H <sub>2</sub> O (1:1)
548	B1a	6-methyl-3-pyridinyl	HCl (1:2)
549	B1a	3-(trifluoromethyl)phenyl	HCl (1:1); mp. 170- 172°C (dec) *
550	B1a	3-methylphenyl	HCl (1:1)
551	B1a	2,3-dichlorophenyl	HCl (1:1); mp. 164- 166°C (dec) *

Co. no.	Ex. no.	Q	Physical data
552	B1a	5-benzo[b]furanyl	HCl (1:1)
553	B1a	3-(methylthio)phenyl	HCl (1:1)
554	B1a	3-hydroxyphenyl	HCl (1:1)
555	B1a	3-methoxyphenyl	HCl (1:1)
556	B1a	3-chlorophenyl	HCl (1:1)
557	B1a	3-(ethoxycarbonyl)phenyl	HCl (1:1)
558	B1a	3-bromophenyl	HCl (1:1)
559	B1a	4-(methylthio)phenyl	HCl (1:1)
560	B1a	4-hydroxyphenyl	HCl (1:1)
561	B1a	4-methoxyphenyl	HCl (1:1)
562	B1a	4-chlorophenyl	HCl (1:1)
563	B1a	4-methylphenyl	HCl (1:1)
564	B1a	4-(trifluoromethyl)phenyl	HCl (1:1)
565	B1a	4-(ethoxycarbonyl)phenyl	HCl (1:1)
566	B1a	4-bromophenyl	
567	B1a	3,4,5-trimethoxyphenyl	
568	B1a	6-(trifluoromethyl)-3-pyridinyl	HCl (1:1); mp 242°C
569	B1a	imidazo[1,2-a]pyridin-6-yl	HCl (1:1); H <sub>2</sub> O (1:3); mp 218°C
570	B1a	4-fluoro-3-methylphenyl	HCl (1:1)


\* = decomposition


Table 10



5


Co. no.	Ex. no.	X	Q	Physical data
126	B1c	CH	2-pyridinyl	HCl (1:1)
127	B1c	CH	4-pyridinyl	HCl (1:2)

Co. no.	Ex. no.	X	Q	Physical data
128	B1a	CH		HCl (1:1)
129	B1c	CH	3-pyridinyl	
3	B1c	N	3-pyridinyl	HBr (1:1)
290	B1b	N	1-naphthalenyl	HBr (1:1)
503	B1a	CH	1-naphthalenyl	HCl (1:1)
571	B1a	CH	6-chloro-3-pyridinyl	HCl (1:1)
572	B1b	N	6-chloro-3-pyridinyl	HBr (1:1)
573	B1a	CH	6-methoxy-3-pyridinyl	
574	B1a	CH	4-methyl-3-pyridinyl	HBr (1:1); H <sub>2</sub> O (1:1)
575	B1b	N	4-methyl-3-pyridinyl	HBr (1:1)
576	B1b	N	6-methoxy-3-pyridinyl	
577	B1a	CH	6-methyl-3-pyridinyl	HCl (1:2); H <sub>2</sub> O (1:2)
578	B1a	CH	6-bromo-3-pyridinyl	HCl (1:1)
579	B1a	CH	2,3-dihydro-5-benzofuranyl	HCl (1:1); mp. 226-228°C (dec) *
580	B1a	CH	5-bromo-3-pyridinyl	HCl (1:1); H <sub>2</sub> O (1:1)
581	B1a	CH	5-chloro-3-pyridinyl	HCl (1:1); H <sub>2</sub> O (1:1)
582	B1a	CH	6-methyl-2-pyridinyl	HCl (1:2); H <sub>2</sub> O (1:1); mp. >250°C
583	B1a	CH	2-methoxy-3-pyridinyl	mp. 222°C
584	B1a	CH	5-(trifluoromethyl)-3-pyridinyl	HCl (1:1); mp. >260°C
585	B1a	CH	5-methyl-2-pyridinyl	HCl (1:1); mp. 230°C
586	B1a	CH	6-(trifluoromethyl)-3-pyridinyl	HCl (1:1); mp. >260°C
587	B1a	CH	6-benzothiazolyl	HCl (1:1); mp. 210-212°C (dec) *
588	B1a	CH	6-hydroxy-3-pyridinyl	HBr (1:2); mp. >260°C
589	B1a	CH	4-methyl-2-pyridinyl	HCl (1:1); H <sub>2</sub> O (1:1)
590	B1a	CH	2,3-dihydro-1,4-benzodioxin-6-yl	HCl (1:1); mp 220°C
591	B1a	CH	1H-indazole-5-yl	HCl (1:1)
592	B1a	CH	6-(methylthio)-3-pyridinyl	HCl (1:1); mp 238°C


Co. no.	Ex. no.	X	Q	Physical data
593	B1a	CH	1 <i>H</i> -benzimidazol-5-yl	HCl (1:1); H <sub>2</sub> O (1:1); mp >260°C
594	B1a	CH	6-ethoxy-3-pyridinyl	HCl (1:2)
595	B1a	CH	1-methyl-1 <i>H</i> -benzimidazol-5-yl	HCl (1:2); H <sub>2</sub> O (1:2); mp 252°C
596	B1a	CH	5-methyl-3-pyridinyl	HCl (1:1); H <sub>2</sub> O (1:1); mp 212°C
597	B1a	CH		HCl (1:2); mp >260°C


\* = decomposition

Table 11



5


Co. no.	Ex. no.	R	Q	Physical data
357	B1b	H		HBr (1:1)
358	B1b	H	1-naphthalenyl	HBr (1:1)
598	B1b	H	3-pyridinyl	HBr (1:2); H <sub>2</sub> O (1:1)
599	B1b	H	6-chloro-3-pyridinyl	HBr (1:1)
600	B1b	H	6-methoxy-3-pyridinyl	
601	B1b	H	4-methyl-3-pyridinyl	HBr (1:2); H <sub>2</sub> O (1:1)
602	B1b	H	5-pyrimidinyl	HBr (1:2)
603	B1b	H	6-bromo-3-pyridinyl	HBr (1:1); H <sub>2</sub> O (1:1)
604	B1b	H	5-chloro-3-pyridinyl	HBr (1:2)
605	B1b	H	2,3-dihydro-5-benzofuranyl	HBr (1:1)
606	B1b	H	5-bromo-3-pyridinyl	HBr (1:2); H <sub>2</sub> O (1:1)
607	B1b	H	6-(trifluoromethyl)-3-pyridinyl	HBr (1:2)

Co. no.	Ex. no.	R	Q	Physical data
608	B1b	H	6-hydroxy-3-pyridinyl	HBr (1:1)
609	B1b	H	2-methoxy-3-pyridinyl	
610	B1b	H	6-benzothiazolyl	HBr (1:1); H <sub>2</sub> O (1:1); mp.>250°C
611	B1b	H	1 <i>H</i> -indazol-5-yl	HBr (1:1); H <sub>2</sub> O (1:1)
612	B1b	H	5-(trifluoromethyl)-3-pyridinyl	HBr (1:1)
613	B1b	H	1 <i>H</i> -benzimidazol-5-yl	HBr (1:2); mp.>260°C
614	B1b	H	2,3-dihydro-1,4-benzodioxin-6-yl	HBr (1:2); mp.>250°C
615	B1b	H	6-ethoxy-3-pyridinyl	HBr (1:2)
616	B1b	H	6-(methylthio)-3-pyridinyl	HBr (1:2); mp >260°C
617	B1b	H	6-methyl-3-pyridinyl	HBr (1:2)
618	B1b	H	1-methyl-1 <i>H</i> -indazol-5-yl	HBr (1:1)
619	B1b	H	1-methyl-1 <i>H</i> -benzimidazol-5-yl	HBr (1:2); mp 246°C
620	B1b	H	5-methyl-3-pyridinyl	HBr (1:2)
621	B1b	5-bromo	3-(trifluoromethyl)phenyl	HB (1:1)
622	B1a	H		HBr (1:2); H <sub>2</sub> O (1:2); mp >260°C
623	B1b	6-CF <sub>3</sub>	3-(trifluoromethyl)phenyl	HBr (1:2); mp 156°C
624	B1b	6-CF <sub>3</sub>	2,3-dichlorophenyl	HBr (1:1); mp 206°C
625	B1b	6-CF <sub>3</sub>	6-methyl-3-pyridinyl	HBr (1:1); mp >260°C
626	B1d	5-CH <sub>3</sub>	3-(trifluoromethyl)phenyl	HBr (1:1)
627	B1d	5-CH <sub>3</sub>	2,3-dichlorophenyl	HBr (1:1)
628	B1b	6-[NH-C(=O)-CH <sub>3</sub> ]	3-(trifluoromethyl)phenyl	mp 248°C
629	B1d	6-CH <sub>3</sub>	3-(trifluoromethyl)phenyl	HBr (1:1)
630	B1b	6-[NH-C(=O)-CH <sub>3</sub> ]	6-methyl-3-pyridinyl	HBr (1:2); H <sub>2</sub> O (1:1); mp >260°C
631	B1d	6-CH <sub>3</sub>	6-methyl-3-pyridinyl	HBr (1:2)
632	B1d	5-CH <sub>3</sub>	6-methyl-3-pyridinyl	HBr (1:2); H <sub>2</sub> O (1:1)
633	B4a	6-NH <sub>2</sub>	6-methyl-3-pyridinyl	HBr (1:2); H <sub>2</sub> O (1:2); mp >260°C

-75-

Co. no.	Ex. no.	R	Q	Physical data
634	B1b	6-[NH-C(=O)-CH <sub>3</sub> ]	2,3-dichlorophenyl	HBr (1:1); mp >260°C
635	B4b	6-NH <sub>2</sub>	2,3-dichlorophenyl	HBr (1:2); mp 236°C
636	B1d	6-CH <sub>3</sub>	2,3-dichlorophenyl	HBr (1:1); H <sub>2</sub> O (1:1)
637	B5	6-NH <sub>2</sub>	3-(trifluoromethyl)phenyl	HBr (1:1); H <sub>2</sub> O (1:1); mp 148°C
638	B1b	6-[C(=O)-NH <sub>2</sub> ]	3-(trifluoromethyl)phenyl	HBr (1:1); ethanolate (1:1)
639	B1b	5-[C(=O)-NH <sub>2</sub> ]	2,3-dichlorophenyl	HBr (1:1); H <sub>2</sub> O (1:2); mp >260°C

Table 12.



5

Co. no.	Ex. no.	Q	Physical data
640	B1b	phenyl	HCl (1:1)
641	B1b	4-methoxyphenyl	
642	B1b	3-pyridinyl	
643	B1b	3-(trifluoromethyl)phenyl	

Table 13 lists both the experimental (column heading "Exper") and theoretical (column heading "Theor") elemental analysis values for carbon (C), hydrogen (H), nitrogen (N) and chlorine (Cl) for the compounds as prepared in the experimental part hereinabove.

10

Table 13

Co. No.	C		H		N		Cl	
	Theor	Exper	Theor	Exper	Theor	Exper	Theor	Exper
18	61.36	60.55	4.88	5.00	11.30	11.09	9.53	9.20
210	47.11	46.45	3.46	3.48	17.17	16.68		
1	51.92	51.74	3.59	3.42	14.25	14.05		
42	42.26	42.28	2.66	2.37	12.32	12.06		

Co. No.	C		H		N		Cl	
	Theor	Exper	Theor	Exper	Theor	Exper	Theor	Exper
61	64.26	63.99	4.79	4.70	16.65	16.43	8.96	9.23
5	48.25	49.23	4.3	4.21	17.58	17.38		
138	44.61	44.59	3.74	3.42	14.86	14.60		
480	47.11	47.08	3.46	3.35	17.17	17.06		
223	50.18	49.96	3.72	3.53	13.77	13.60		
239	56.9	56.97	4.21	3.96	15.61	15.32		
291	57.64	57.10	3.33	3.02	29.41	29.11		
299	58.3	58.36	4.08	3.94	15.11	14.94		
93	54.70	54.49	3.51	3.30	7.50	7.39		
113	62.52	62.06	4.26	4.19	22.78	22.63		
125	43.08	43.79	2.99	2.63	15.96	15.63	0	0.15
140	36.22	39.32	3.47	3.69	12.07	12.62		
511	42.97	42.78	3.37	3.16	13.36	13.09		
517	52.17	51.94	2.81	2.61	17.38	16.90		
518	48.49	48.08	2.91	2.66	8.08	7.94		
132	57.13	56.27	3.42	3.27	28.55	28.37		
133	45.93	45.55	3.6	3.22	17.85	17.40		
150	65.35	65.46	6.45	6.45	17.93	18.08		
218	61.15	60.49	6.04	6.24	12.59	12.16		
602	34.55	33.85	2.66	2.98	16.79	16.28		
525	50.18	49.67	2.59	2.64	9	8.54	8.05	16.95
526	55.54	54.94	3.73	3.56	25.91	24.99		
527	40.38	39.09	3.39	3.16	18.11	17.26		
149	40.68	41.01	3.17	2.82	13.56	13.65		
546	42.17	42.17	2.45	2.20	18.91	18.60		
569	38.71	39.96	3.9	3.52	18.06	18.18		
579	58.3	58.25	4.08	3.75	15.11	14.93		
587	52.91	52.70	3.13	3.02	18.15	18.23		
590	55.89	55.67	3.91	3.94	14.48	14.42		
591	55.36	54.32	3.55	3.47	22.78	22.26		
593	52.78	54.64	3.91	4.10	21.72	22.90	9.42	9.71
605	51.07	51.01	3.75	3.46	11.17	10.94		
623	34.87	36.06	2.01	2.08	7.62	7.75		
628	53.96	53.23	3.46	3.17	14.81	14.50		
638	44	43.13	3.69	3.26	11.4	11.27		

Co.	C		H		N		Cl	
No.	Theor	Exper	Theor	Exper	Theor	Exper	Theor	Exper
128	54.77	54.65	3.51	3.22	15.03	14.81		
634	41.76	41.60	2.85	2.63	12.18	11.85		
643	51.28	51.51	2.96	2.81	7.47	7.42		

### C. Pharmacological example

#### Example C.1 : *in vitro* inhibition of TNF- $\alpha$ production in human blood

##### *Human whole blood stimulation*

- 5 Peripheral blood from healthy male donors was drawn into heparinized syringes (12.5 U heparin/ml). Blood samples were three-fold diluted in RPMI 1640 medium (Life Technologies, Belgium) supplemented with 2 mM L-glutamine, 100 U/ml penicillin and 100  $\mu$ g/ml streptomycin, and 300  $\mu$ l fractions were distributed in 24-well multidisc plates (Nunc, Roskilde, Denmark). Blood samples were preincubated (60
- 10 minutes at 37°C) in a humidified 6% CO<sub>2</sub>-atmosphere with 100  $\mu$ l of drug solvent (final concentration 0.02% dimethylsulfoxide in RPMI 1640) or with 100  $\mu$ l of an appropriate dose of test compound before being stimulated by the addition of 100  $\mu$ l of lipopolysaccharide at a final concentration of 100 ng/ml. After 6 hours, cell-free supernatant fluids were collected by centrifugation and stored at -20°C until tested for
- 15 the presence of TNF- $\alpha$ .

#### Example C.2 : *in vitro* inhibition of IL-12 production in human blood

##### *Human whole blood stimulation*

- Peripheral blood from healthy male donors was drawn into heparinized syringes (12.5 U heparin/ml). Blood samples were three-fold diluted in RPMI 1640 medium (Life Technologies, Belgium) supplemented with 2 mM L-glutamine, 100 U/ml penicillin and 100  $\mu$ g/ml streptomycin, and 300  $\mu$ l fractions were distributed in 24-well multidisc plates (Nunc, Roskilde, Denmark). Blood samples were preincubated (60
- 20 minutes at 37°C) in a humidified 6% CO<sub>2</sub>-atmosphere with 100  $\mu$ l of drug solvent (final concentration 0.02% dimethylsulfoxide in RPMI 1640) or with 100  $\mu$ l of an appropriate dose of test compound before being stimulated by the addition of 100  $\mu$ l of lipopolysaccharide at a final concentration of 100 ng/ml. After 24 hours, cell-free supernatant fluids were collected by centrifugation and stored at -20°C until tested for
- 25 the presence of IL-12.

Example C.3 : cytokine measurements


- Cytokine protein concentrations were determined by sandwich ELISA as described in Van Wauwe et al. (1996, Inflamm Res, 45, 357-363). Murine monoclonals used as capture antibodies to human cytokines were obtained from R&D Systems (Abingdon, United Kingdom) and code named MAB210 and MAB611 for TNF- $\alpha$  and IL-12 respectively. Biotinylated goat polyclonal antibodies used to detect human cytokines were from R&D Systems (BAF210, BAF219). Cytokine levels were calculated from standard curves using recombinant cytokines supplied by R&D Systems.
- 10 Table 14 lists the percentage inhibition of TNF- $\alpha$  and IL-12 production (column “%inh”) at a test dose of  $1 \times 10^{-6}$  and  $1 \times 10^{-7}$  M for the compounds of the present invention.

Table 14

Comp. No	% inhib. TNF- $\alpha$		% inhib. IL-12 (p40)	
	$1 \times 10^{-6}$ M	$1 \times 10^{-7}$ M	$1 \times 10^{-6}$ M	$1 \times 10^{-7}$ M
9	37	39	49	53
140	46	44	56	63
74	56	48	70	67
81	51	47	65	67
82	53	51	73	68
100	43	41	53	51
101	54	53	62	65
31	55	49	66	68
39	53	59	64	71
476	58	53	75	71
45	49	48	64	65
166	48	37	62	55
410	39	43	53	58
115	58	53	75	67
119	49	49	62	62
286	50	48	60	63
573	53	45	67	61
526	45	45	66	69
577	50	49	77	71
527	37	43	61	66
549	50	47	74	71
551	44	40	71	71
579	49	50	72	75
584	53	49	75	68
587	56	56	79	75
517	61	57	74	68
643	64	59	75	72
518	38	44	62	59
466	57	49	95	86
509	46	54	64	68

Claims

1. Use of a compound for the manufacture of a medicament for the prevention or the treatment of diseases mediated through cytokines, wherein the compound is a compound of formula




a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof,  
wherein

10 Q is C<sub>3-6</sub>cycloalkyl, phenyl, naphthyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxy; azido; amino; mono- or di(C<sub>1-6</sub>alkyl)amino; C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>2-6</sub>alkynyl; C<sub>3-6</sub>cycloalkyl; C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, amino, mono- or di(C<sub>1-4</sub>alkyl)amino;  
15 C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylthio; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl; arylC<sub>1-6</sub>alkyloxy; aryloxy; polyhaloC<sub>1-6</sub>alkyl; polyhalo-C<sub>1-6</sub>alkyloxy; polyhalo-C<sub>1-6</sub>alkylcarbonyl; C<sub>1-4</sub>alkyl-S(=O)<sub>n</sub>- or R<sup>1</sup>HN-S(=O)<sub>n</sub>-;

or

Q is a radical of formula



wherein X and Y each independently are O, NR<sup>3</sup>, CH<sub>2</sub> or S, with R<sup>3</sup> being hydrogen or C<sub>1-4</sub>alkyl;


q is an integer with value 1 to 4;

Z is O or NR<sup>4</sup> with R<sup>4</sup> being hydrogen or C<sub>1-4</sub>alkyl;

25 r is an integer with value 1 to 3;

n is an integer with value 1 or 2;

R<sup>1</sup> represents hydrogen, or a radical of formula



with A being O, S or a bivalent radical of formula -CR<sup>2a</sup>=N- with CR<sup>2a</sup> attached to N of formula (a-1); and

R<sup>2a</sup> being hydrogen, C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy;

L is phenyl, optionally substituted with up to 4 substituents each independently being selected from halo, hydroxy, amino, cyano, carboxyl, mono- or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono- or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-; or


L is Het;

- Het is (i) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each independently being selected where possible from N, S or O;
- (ii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each independently being selected where possible from N, S or O and being fused through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered ring, which contains, apart from the atoms in common with the first ring, only carbon atoms; the latter ring may be unsaturated, partially unsaturated or saturated;
- (iii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and at least one heteroatom and being fused through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered heterocyclic ring, which contains, apart from the atoms in common with the first ring, at least one heteroatom; the latter ring may be unsaturated, partially unsaturated or saturated; said bicyclic ring system contains in total from 2 up to 6 heteroatoms, each independently being selected where possible from N, S or O;

wherein Het being a monocyclic ring system may optionally be substituted with up to 4 substituents, and wherein Het being a bicyclic ring system may optionally be substituted with up to 6 substituents, said substituents each independently being selected from halo, hydroxy, amino, cyano, carboxyl, mono- or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono- or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-;

aryl is phenyl, optionally substituted with up to five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylthio, cyano, nitro, amino, mono- or di(C<sub>1-6</sub>alkyl)amino.

- 5 2. Use of a compound as claimed in claim 1 wherein L is phenyl, optionally substituted with up to 4 substituents each independently being selected from halo, hydroxy, amino, mono or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy; or L is Het, optionally substituted with up to 4 substituents when Het is a monocyclic ring, or up to 6 substituents when Het is a bicyclic ring, said substituents each  
 10 independently being selected from halo, hydroxy, amino, mono or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy; and wherein aryl is phenyl, optionally substituted with up to five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylthio, cyano, nitro, amino, mono- or di(C<sub>1-6</sub>alkyl)amino; and wherein Q is phenyl, naphthyl, pyridyl,  
 15 pyrimidinyl, pyrazinyl or pyridazinyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxy; azido; amino; mono- or di(C<sub>1-6</sub>alkyl)amino; C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>2-6</sub>alkynyl; C<sub>3-6</sub>cycloalkyl; C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, amino, mono- or di(C<sub>1-4</sub>alkyl)amino; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylthio; C<sub>1-6</sub>alkylcarbonyl;  
 20 C<sub>1-6</sub>alkyloxycarbonyl; arylC<sub>1-6</sub>alkyloxy; aryloxy; polyhaloC<sub>1-6</sub>alkyl; polyhalo-C<sub>1-6</sub>alkyloxy; polyhaloC<sub>1-6</sub>alkylcarbonyl; C<sub>1-4</sub>alkyl-S(=O)<sub>n</sub>- or R<sup>1</sup>HN-S(=O)<sub>n</sub>-; or Q is a radical of formula




wherein X and Y each independently are O, NR<sup>3</sup>, CH<sub>2</sub> or S, with R<sup>3</sup> being

- 25 hydrogen or C<sub>1-4</sub>alkyl;  
 q is an integer with value 1 to 4;  
 Z is O or NR<sup>4</sup> with R<sup>4</sup> being hydrogen or C<sub>1-4</sub>alkyl;  
 r is an integer with value 1 to 3;

- 30 3. Use of a compound as claimed in claim 1 or 2 provided that L is other than optionally substituted phenyl and provided that when L is Het then Het is other than optionally substituted isothiazolyl, 2-pyridyl, benzthiazolyl, benzoxazinyl and benzoxazinonyl.

4. Use of a compound as defined in any one of claims 1 to 3 for the manufacture of a medicament for the prevention or the treatment of diseases mediated through activation of the adenosine A<sub>3</sub> receptor.
5. Use according to any one of claims 1 to 3 wherein the diseases mediated through cytokines are TNF- $\alpha$  (Tumor Necrosis Factor-alpha) and IL-12 (Interleukin 12) mediated diseases.
6. Use according to any one of claims 1, 2, 3 and 5 wherein the diseases are inflammatory or autoimmune diseases.
7. Use according to claim 6 wherein the inflammatory or autoimmune diseases are rheumatoid arthritis, Crohn's disease, irritable bowel disease or colitis.
8. A compound of formula




a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof,  
wherein

- Q is C<sub>3-6</sub>cycloalkyl, phenyl, naphthyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzthiazolyl, benzoxazolyl, benzimidazolyl, indazolyl, or imidazopyridyl, each of said rings optionally being substituted with up to three substituents each independently selected from halo; hydroxy; cyano; carboxy; azido; amino; mono- or di(C<sub>1-6</sub>alkyl)amino; C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>2-6</sub>alkynyl; C<sub>3-6</sub>cycloalkyl; C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, amino, mono- or di(C<sub>1-4</sub>alkyl)amino; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylthio; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl; arylC<sub>1-6</sub>alkyloxy; aryloxy; polyhaloC<sub>1-6</sub>alkyl; polyhalo-C<sub>1-6</sub>alkyloxy; polyhalo-C<sub>1-6</sub>alkylcarbonyl; C<sub>1-4</sub>alkyl-S(=O)<sub>n</sub>- or R<sup>1</sup>HN-S(=O)<sub>n</sub>-;

or

- Q is a radical of formula



wherein X and Y each independently are O, NR<sup>3</sup>, CH<sub>2</sub> or S, with R<sup>3</sup> being hydrogen or C<sub>1-4</sub>alkyl;

-84-


q is an integer with value 1 to 4;

Z is O or NR<sup>4</sup> with R<sup>4</sup> being hydrogen or C<sub>1-4</sub>alkyl;

r is an integer with value 1 to 3;

n is an integer with value 1 or 2;

- 5 R<sup>1</sup> represents hydrogen, or a radical of formula



with A being O, S or a bivalent radical of formula -CR<sup>2a</sup>=N- with CR<sup>2a</sup> attached to N of formula (a-1); and

R<sup>2a</sup> being hydrogen, C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy;

- 10 L is 3-halophenyl, optionally substituted with 1, 2 or 3 substituents each independently being selected from halo, hydroxy, amino, cyano, carboxyl, mono- or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono- or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxy-carbonyl, C<sub>1-6</sub>alkyl-carbonyloxy, aminocarbonyl, 15 mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-; or

L is Het;

Het is (i) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each 20 independently being selected where possible from N, S or O;

(ii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and containing 1, 2, 3 or 4 heteroatoms each 25 independently being selected where possible from N, S or O and being fused through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered ring, which contains, apart from the atoms in common with the first ring, only carbon atoms; the latter ring may be unsaturated, partially unsaturated or saturated;


(iii) an optionally substituted five- or six-membered heterocyclic ring containing at least two double bonds and at least one heteroatom and being fused 30 through 2 carbon atoms, 2 nitrogen atoms or 1 carbon and 1 nitrogen atom with another optionally substituted five- or six-membered heterocyclic ring, which contains, apart from the atoms in common with the first ring, at least one heteroatom; the latter ring may be unsaturated, partially unsaturated or saturated; said bicyclic ring system contains in total from 2

up to 6 heteroatoms, each independently being selected where possible from N, S or O;

- wherein Het being a monocyclic ring system may optionally be substituted with up to 4 substituents, and wherein Het being a bicyclic ring system may optionally be substituted with up to 6 substituents, said substituents each independently being selected from halo, hydroxy, amino, cyano, carboxyl, mono-or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted with hydroxy or C<sub>1-4</sub>alkyloxy or amino or mono-or di(C<sub>1-4</sub>alkyl)amino, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyloxy, aminocarbonyl, mono-or di(C<sub>1-6</sub>alkyl)aminocarbonyl, C<sub>1-6</sub>alkyl-C(=O)-NH-, C<sub>1-6</sub>alkyloxy-C(=O)-NH-, H<sub>2</sub>N-C(=O)-NH- or mono- or di(C<sub>1-4</sub>alkyl)amino-C(=O)-NH-; aryl is phenyl, optionally substituted with up to five substituents each independently selected from halo, hydroxy, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylthio, cyano, nitro, amino, mono-or di(C<sub>1-6</sub>alkyl)amino.
- provided that
- Het is other than optionally substituted isothiazolyl, 2-pyridyl, benzthiazolyl, benzoxazinyl and benzoxazinonyl;
  - when Q is phenyl substituted with hydroxy or C<sub>1-6</sub>alkyloxy and carboxy or C<sub>1-6</sub>alkyloxycarbonyl then Het is other than 3-pyridyl or 4-pyridyl;
  - when Q is phenyl then Het is other than 2-thienyl, 2-furanyl, 5-bromo-2-benzofuranyl, 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl, 2-benzofuranyl, 5-chloro-2-benzimidazolyl, 2-benzimidazolyl, 3-pyridyl, 4-pyridyl, 6-methyl-thiazolo [3,2-b] [1,2,4]triazol-5-yl, 2,6-dimethyl-thiazolo [3,2-b] [1,2,4]triazol-5-yl or 5,6-dihydro-4,5-dimethyl-2(H)-3-pyranonyl;
  - when Q is 2-methyl-phenyl then Het is other than 2-thienyl, 2-benzofuranyl or 3-pyridyl;
  - when Q is 4-methoxy-phenyl then Het is other than 2-furanyl, 2-pyrazinyl, 3-pyridyl, 4-pyridyl, 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl, 1,2-dihydro-6-ethyl-2-oxo-3-cyano-5-pyridyl, 4-(dimethylamino)-1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl, 1,2-dihydro-4-methoxy-6-methyl-2-oxo-3-cyano-5-pyridyl or 3-amino-6-methyl-2(1H)-5-pyridinonyl;
  - when Q is 2-methoxy-phenyl then Het is other than 2-pyrazinyl, 5-chloro-2-benzimidazolyl, or 3-pyridyl;
  - when Q is 4-chloro-phenyl then Het is other than 2-furanyl, 2-thienyl, 5-chloro-2-benzimidazolyl, 2-pyrazinyl, 3-pyridyl, 4-pyridyl or 5,6-dihydro-4,5-dimethyl-2(H)-3-pyranonyl;

- when Q is 3-chloro-phenyl then Het is other than 2-thienyl, 3-pyridyl or 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl;
  - when Q is 2-chloro-phenyl then Het is other than 2-thienyl;
  - when Q is 3-methyl-phenyl then Het is other than 2-thienyl or 3-pyridyl;
  - 5 - when Q is 2,3-dichloro-phenyl then Het is other than 3-pyridyl;
  - when Q is 2-ethoxy-phenyl or 3-methoxy-phenyl then Het is other than 2-pyrazinyl;
  - when Q is 4-bromo-phenyl then Het is other than 2-thienyl, or 5-chloro-2-benzimidazolyl;
  - when Q is 4-fluoro-phenyl then Het is other than 4-pyridyl;
  - 10 - when Q is 1-naphthyl then Het is other than 2-thienyl, or 3-pyridyl;
  - when Q is 4-methyl-phenyl then Het is other than 2-furanyl, 2-thienyl, 3-pyridyl, 2-pyrazinyl or 5,6-dihydro-4,5-dimethyl-2(H)-3-pyranonyl;
  - when Q is 4-ethoxy-phenyl then Het is other than 2-pyrazinyl;
  - when Q is 2-naphthyl, 2-carboxy-phenyl, 3-carboxy-phenyl, 4-carboxy-phenyl,
  - 15 4-amino-phenyl or 3-chloro-2,6-dinitro-4-trifluoromethyl-phenyl then Het is other than 2-thienyl;
  - when Q is 4-benzenesulfonamide then Het is other than 2-furanyl, or 1,2,3,4-tetrahydro-2,4-dioxo-5-pyrimidinyl;
  - when Q is N-methyl-4-benzenesulfonamide then Het is other than 3-thienyl;
  - 20 - when Q is N-butyl-4-benzenesulfonamide then Het is other than 2-furanyl;
  - when Q is 2-pyridyl then Het is other than 2-pyrazinyl;
  - when Q is 3-pyridyl then Het is other than 2-thienyl, 3-pyridyl, 4-pyridyl or 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl;
  - when Q is 2,4-dichloro-phenyl then Het is other than 2-pyrazinyl, or 4-pyridyl;
  - 25 when Q is 4-pyridyl then Het is other than 3-quinoliny or 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl;
  - when Q is 2,4-dimethoxyphenyl, 3,4-dimethoxyphenyl, 4-hydroxyphenyl, 4-methylthiophenyl, or 4-methylsulfinylphenyl then Het is other than 1,2-dihydro-6-methyl-2-oxo-3-cyano-5-pyridyl.
- 30
9. A compound as claimed in claim 8 wherein L is Het and Het, being a monocyclic ring system, may optionally be substituted with up to 4 substituents, and Het, being a bicyclic ring system, may optionally be substituted with up to 6 substituents, said substituents each independently being selected from halo, hydroxy, amino, mono or
- 35 di(C<sub>1-4</sub>alkyl)amino, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy; and wherein Q is phenyl, naphthyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl, each of said rings optionally being substituted with up to three substituents each independently selected

- from halo; hydroxy; cyano; carboxy; amino; mono- or di(C<sub>1-6</sub>alkyl)amino; C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>2-6</sub>alkynyl; C<sub>3-6</sub>cycloalkyl; C<sub>1-6</sub>alkyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, amino, mono- or di(C<sub>1-4</sub>alkyl)amino; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylthio; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl; arylC<sub>1-6</sub>alkyloxy; aryloxy; polyhaloC<sub>1-6</sub>alkyl; polyhaloC<sub>1-6</sub>alkyloxy; polyhaloC<sub>1-6</sub>alkylcarbonyl; C<sub>1-4</sub>alkyl-S(=O)<sub>n</sub>- or R<sup>1</sup>HN-S(=O)<sub>n</sub>-; or Q is a radical of formula



wherein X and Y each independently are O, NR<sup>3</sup>, CH<sub>2</sub> or S, with R<sup>3</sup> being hydrogen or C<sub>1-4</sub>alkyl;

q is an integer with value 1 to 4;

Z is O or NR<sup>4</sup> with R<sup>4</sup> being hydrogen or C<sub>1-4</sub>alkyl;

r is an integer with value 1 to 3;

10. A compound as claimed in claim 8 wherein L is imidazolyl, imidazothiazolyl, pyrimidinyl, thienyl, thiazolyl, furanyl, 3-pyridyl, 4-pyridyl, pyrazolyl, indolyl, indazolyl, quinolinyl, benzofuranyl, pyrrolopyridyl, imidazopyridyl, imidazopyrazinyl, imidazopyrimidinyl, imidazopyridazinyl, pyrazolopyridyl, with each heterocycle optionally substituted with one, two, three or four substituents selected from halo, amino, C<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl, aminocarbonyl or C<sub>1-6</sub>alkyl-C(=O)-NH-.
11. A compound as claimed in any one of claims 8 to 10 wherein L is indolyl, 3-imidazo[1,2-a]pyridyl, 3-imidazo[1,5-a]pyridyl, 3-pyridyl, quinolinyl, imidazopyrimidinyl, imidazopyrazinyl, imidazothiazolyl, 5-pyrimidinyl, furanyl, thiazolyl, imidazolyl, pyrrolopyridyl, pyrazolopyridyl.
12. A compound as claimed in any one of claims 8 to 11 wherein L is 3-imidazo[1,2-a]pyridyl, 3-imidazo[1,5-a]pyridyl, imidazothiazolyl, 3-pyridyl or pyrrolopyridyl.
13. A compound as claimed in claim 8 wherein L is 3-fluorophenyl or 3,5-difluorophenyl.

14. A compound as claimed in any one of claims 8, 10, 11, 12 or 13 wherein Q is 3-pyridyl, 4-pyridyl, naphthalenyl, C<sub>3-6</sub>cycloalkyl, phenyl, 1,3-benzodioxolyl, 2,3-dihydro-benzofuranyl, 2,3-dihydro-1,4-benzodioxinyl, benzthiazolyl, indazolyl, benzimidazolyl or imidazopyridyl.

5

15. A compound as claimed in any one of claims 8, 10, 11, 12, 13 or 14 wherein Q is phenyl, 3-pyridyl, 4-pyridyl, benzthiazolyl or imidazopyridyl, each of said rings being optionally substituted with up to three substituents selected from halo, cyano, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy or polyhaloC<sub>1-6</sub>alkyl.

10

16. A compound as claimed in claim 8 wherein the compound is selected from

- 2-thiazolamine, 4-imidazo[1,2-a]pyridin-3-yl-N-[3-(trifluoromethyl)phenyl];
- 2-thiazolamine, 4-imidazo[1,2-a]pyridin-3-yl-N-[4-(trifluoromethyl)phenyl];
- 2-thiazolamine, 4-(3-pyridinyl)-N-[3-(trifluoromethyl)phenyl];
- 15 2-thiazolamine, N-(3-chlorophenyl)-4-imidazo[1,2-a]pyridin-3-yl;
- 2-thiazolamine, 4-imidazo[1,2-a]pyridin-3-yl-N-(3-methylphenyl);
- 2-thiazolamine, 4-imidazo[1,2-a]pyridin-3-yl-N-[3-(methylthio)phenyl];
- 2-thiazolamine, N-(4-chlorophenyl)-4-imidazo[1,2-a]pyridin-3-yl;
- 2-thiazolamine, N-(3-bromophenyl)-4-imidazo[1,2-a]pyridin-3-yl;
- 20 2-thiazolamine, N-(2,3-dichlorophenyl)-4-imidazo[1,2-a]pyridin-3-yl;
- 2-thiazolamine, N-(2,3-dichlorophenyl)-4-(1H-pyrrolo[2,3-b]pyridin-3-yl);
- 2-thiazolamine, N-(4-bromophenyl)-4-imidazo[1,2-a]pyridin-3-yl;
- 2-thiazolamine, N-(2,3-dichlorophenyl)-4-imidazo[1,5-a]pyridin-3-yl;
- 2-thiazolamine, 4-imidazo[2,1-b]thiazol-5-yl-N-[3-(trifluoromethyl)phenyl];
- 25 2-thiazolamine, N-(2,3-dichlorophenyl)-4-imidazo[2,1-b]thiazol-5-yl;
- 2-thiazolamine, 4-(3-pyridinyl)-N-(3-methyl-4-fluorophenyl);
- 2-thiazolamine, 4-imidazo[1,2-a]pyridin-3-yl-N-(3-methyl-4-fluorophenyl);
- a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and stereochemically isomeric form thereof.

30

17. A compound as claimed in claim 8 or 13 wherein the compound is selected from


- 2-thiazolamine, 4-(3-fluorophenyl)-*N*-phenyl;
- 2-thiazolamine, 4-(3-fluorophenyl)-*N*-[4-methoxyphenyl];
- 2-thiazolamine, 4-(3-fluorophenyl)-*N*-[4-(trifluoromethyl)phenyl]; and
- 35 2-thiazolamine, 4-(3-fluorophenyl)-*N*-[3-pyridyl];
- a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof.

18. A compound as claimed in any one of claims 8 to 17 for use as a medicine.

19. A pharmaceutical composition comprising a pharmaceutically acceptable carrier  
5 and as active ingredient a therapeutically effective amount of a compound as  
claimed in any one of claims 8 to 17.

20. A process of preparing a composition as claimed in claim 19 characterized in that a  
pharmaceutically acceptable carrier is intimately mixed with a therapeutically  
10 effective amount of a compound as claimed in any one of claims 8 to 17.


21. A process of preparing a compound as claimed in claim 8 characterized by  
a) reacting an intermediate of formula (II) or formula (III) or reacting an  
intermediate of formula (II) and (III) with an intermediate of formula (IV)



15

with L and Q defined as in claim 8 and W<sub>1</sub> being a suitable leaving group, in a  
suitable reaction-inert solvent,


b) reacting an intermediate of formula (II-a) with an intermediate of formula (IV)



20

with Q defined as in claim 8, H<sub>2</sub>N-L<sub>1</sub> being defined as L according to claim 1  
provided that L is substituted with NH<sub>2</sub>, and W<sub>1</sub> being a suitable leaving group, in a  
suitable reaction-inert solvent and in the presence of a suitable acid;

c) reducing an intermediate of formula (I-b-interm.)



with L defined as in claim 8,  $\text{Q}_1-\text{NH}_2$  being defined as Q according to claim 8 provided that Q is substituted with  $\text{NH}_2$ , in a suitable solvent and in the presence of a suitable reducing agent;

and, if desired, converting compounds of formula (I) into each other following art-known transformations, and further, if desired, converting the compounds of formula (I), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or into a therapeutically active non-toxic base addition salt by treatment with a base, or conversely, converting the acid addition salt form into the free base by treatment with alkali, or converting the base addition salt into the free acid by treatment with acid; and, if desired, preparing stereochemically isomeric forms, quaternary amines or *N*-oxide forms thereof.

22. A product containing (a) a compound as defined in claim 8, and (b) another anti-inflammatory or immunosuppressive compound, as a combined preparation for simultaneous, separate or sequential use in the treatment of inflammatory or autoimmune diseases.

23. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and as active ingredients (a) a compound as defined in claim 8, and (b) another anti-inflammatory or immunosuppressive compound.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/01841

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D417/04 A61K31/425 A61P29/00 C07D471/04 C07D417/12  
C07D277/42 C07D487/04 C07D513/04 C07D417/14

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

PAJ, EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,Y	JP 62 108859 A (OTSUKA PHARMACEUT FACTORY INC) 20 May 1987 (1987-05-20) example 13 & CHEMICAL ABSTRACTS, vol. 107, no. 21, 23 November 1987 (1987-11-23) Columbus, Ohio, US; abstract no. 197776, abstract ---	1-23
X,Y	JP 62 067023 A (OTSUKA PHARMACEUT FACTORY INC) 26 March 1987 (1987-03-26) example 8 & CHEMICAL ABSTRACTS, vol. 107, no. 8, 24 August 1987 (1987-08-24) Columbus, Ohio, US; abstract no. 64875, abstract --- -/--	1-23

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*8\* document member of the same patent family

Date of the actual completion of the international search

19 June 2001

Date of mailing of the international search report

05/07/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Steendijk, M

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/01841

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,Y	JP 62 067022 A (OTSUKA PHARMACEUT FACTORY INC) 26 March 1987 (1987-03-26) example 8 & CHEMICAL ABSTRACTS, vol. 107, no. 8, 24 August 1987 (1987-08-24) Columbus, Ohio, US; abstract no. 64874, abstract ---	1-23
X,Y	JP 04 154773 A (GREEN CROSS CORP:THE) 27 May 1992 (1992-05-27) page 1 & CHEMICAL ABSTRACTS, vol. 117, no. 21, 23 November 1992 (1992-11-23) Columbus, Ohio, US; abstract no. 212486, abstract ---	1-23
X,Y	CHEMICAL ABSTRACTS, vol. 83, no. 23, 8 December 1975 (1975-12-08) Columbus, Ohio, US; abstract no. 193158, SAWHNEY ET AL.: "Synthesis and antiinflammatory activity of ..." XP002143922 & J.Indian.Chem.Soc.1975,52(6),561-562 abstract ---	1-23
Y	CHEMICAL ABSTRACTS, vol. 111, no. 3, 17 July 1989 (1989-07-17) Columbus, Ohio, US; abstract no. 23457, SAWHNEY ET AL.: "Synthesis of some 2-heterocyclylpheothiazines ..." XP002143923 & J.Indian.Chem.Soc.,1988,65(9),643-647 abstract ---	1-23
Y	CHEMICAL ABSTRACTS, vol. 108, no. 25, 20 June 1988 (1988-06-20) Columbus, Ohio, US; abstract no. 221648, SASTRY ET AL.: "Synthesis and biological activity..." XP002143924 & Indian J.Chem.,Sect.B, 1987,26B(7),662-665 abstract ---	1-23
Y	GB 2 022 085 A (PFIZER) 12 December 1979 (1979-12-12) claim 1 --- -/--	1-23

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/01841

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	JP 55 133366 A (OTSUKA PHARMACEUT FACTORY INC) 17 October 1980 (1980-10-17) page 1 & CHEMICAL ABSTRACTS, vol. 94, no. 17, 27 April 1981 (1981-04-27) Columbus, Ohio, US; abstract no. 139797, abstract ---	1-23
P,X	WO 00 33837 A (SMITHKLINE BEECHAM CORP ;LAGO MARIA A (US)) 15 June 2000 (2000-06-15) the whole document ---	1-14
P,X	WO 00 17175 A (BEMIS GUY ;VERTEX PHARMA (US); COCHRAN JOHN (US); SALITURO FRANCES) 30 March 2000 (2000-03-30) the whole document ---	8-15
X,Y	CHEMICAL ABSTRACTS, vol. 82, no. 5, 3 February 1975 (1975-02-03) Columbus, Ohio, US; abstract no. 31289, SAWHNEY, S. N. ET AL: "Benzothiazole derivatives. III. Synthesis and antiinflammatory activity of some 2-(2-amino-4-thiazolyl)benzothiazoles" XP002151288 abstract & J. INDIAN CHEM. SOC. (1974), 51(5), 566-8, ---	1-23
X,Y	CHEMICAL ABSTRACTS, vol. 90, no. 9, 26 February 1979 (1979-02-26) Columbus, Ohio, US; abstract no. 72098, ARYA, V. P. ET AL: "Psychoactive agents: part VIII. Isothiazoles: part VIII. Synthesis an biological activity of 4-(2-amino-4-thiazolyl)isothiazoles" XP002151289 abstract & INDIAN J. CHEM., SECT. B (1978), 16B(5), 402-4, --- -/--	1-23

# INTERNATIONAL SEARCH REPORT

Inter. Appl. No.

PCT/EP 01/01841

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHEMICAL ABSTRACTS, vol. 103, no. 7, 19 August 1985 (1985-08-19) Columbus, Ohio, US; abstract no. 53986, HANMANTGAD, S. S. ET AL: "Biomimetic thiazolyl coumarins" XP002151290 abstract & NATL. ACAD. SCI. LETT. (INDIA) (1984), 7(3), 77-8,	1-23
Y	--- CHEMICAL ABSTRACTS, vol. 88, no. 15, 10 April 1978 (1978-04-10) Columbus, Ohio, US; abstract no. 105204, SAWHNEY, S. N. ET AL: "Thiazole derivatives: Part II. Synthesis of some 2,4'- and 4,4-bithiazolyl derivatives as potential anti-inflammatory agents" XP002151291 abstract & INDIAN J. CHEM., SECT. B (1977), 15(8), 727-30,	1-23
Y	--- CHEMICAL ABSTRACTS, vol. 94, no. 7, 16 February 1981 (1981-02-16) Columbus, Ohio, US; abstract no. 47195, UPADHYA, KRISHNA G. ET AL: "Synthesis and antiinflammatory activity of 3-substituted 4-(4'-thiazolyl)sydnones" XP002151292 abstract & ARCH. PHARM. (WEINHEIM, GER.) (1980), 313(8), 684-8,	1-23
X	--- CHEMICAL ABSTRACTS, vol. 89, no. 23, 4 December 1978 (1978-12-04) Columbus, Ohio, US; abstract no. 197384, THAKAR ET AL.: "Synthesis of 2-(substituted anilino)4-(substituted phenyl)thiazoles" XP002170057 & J. Pharm. Sci. 1978, 67(4), 587-589 abstract	4, 8
Y	--- WO 99 21555 A (KANZAKI NAUYUKI ;KIMURA HIROYUKI (JP); OHKAWA SHIGENORI (JP); TAKE) 6 May 1999 (1999-05-06) cited in the application claim 1 --- -/--	1-23

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/01841

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	<p>DATABASE CHEMCATS 'Online!                      Accession No. 1998:629695,                      3 April 2000 (2000-04-03)                      XP002170058                      RN: 256471-44-0</p> <p>-----</p>	17

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/ EP/ . . 01/01841

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

SEE ANNEXE

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 8-15, 18-23

Present claims 8-15, 18-23 relate to an extremely large number of possible compounds, products and methods. In fact, the claims contain so many options, variables, possible permutations combined with provisos that a lack of clarity and conciseness within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible. Consequently, the search has been carried out for those parts of the application which do allow a meaningful search, namely the use of the relevant compounds as defined in claims 1 and 4 and compounds as defined in claims 16 and 17.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. .onal Application No

PCT/EP 01/01841

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 62108859 A	20-05-1987	JP 1746098 C JP 4025947 B	25-03-1993 06-05-1992
JP 62067023 A	26-03-1987	JP 1817342 C JP 5024889 B	18-01-1994 09-04-1993
JP 62067022 A	26-03-1987	NONE	
JP 04154773 A	27-05-1992	NONE	
GB 2022085 A	12-12-1979	AR 226285 A AT 373248 B AT 401679 A AU 511242 B AU 4763479 A BE 876732 A CA 1117949 A CH 639653 A CS 216927 B DD 144055 A DE 2922523 A DK 177679 A, B, EG 14354 A ES 481220 A FI 791754 A, B, FR 2427333 A GR 73142 A HK 66487 A HU 180045 B IE 48426 B IL 57450 A IT 1121238 B JP 1261864 C JP 54160369 A JP 59036988 B KE 3459 A LU 81349 A MY 31885 A NL 7904337 A, B, NO 791831 A, B, NZ 190623 A PH 17020 A PL 216030 A PT 69718 A SE 438333 B SE 7904798 A SG 56184 G SU 843746 A US 4307106 A YU 128879 A ZA 7902729 A	30-06-1982 27-12-1983 15-05-1983 07-08-1980 06-12-1979 03-12-1979 09-02-1982 30-11-1983 31-12-1982 24-09-1980 06-12-1979 03-12-1979 30-06-1984 01-09-1980 26-11-1979 28-12-1979 09-02-1984 25-09-1987 28-01-1983 23-01-1985 30-11-1982 26-03-1986 25-04-1985 19-12-1979 06-09-1984 12-10-1984 07-12-1979 31-12-1985 04-12-1979 04-12-1979 06-07-1984 11-05-1984 10-03-1980 01-07-1979 15-04-1985 03-12-1979 08-03-1985 30-06-1981 22-12-1981 31-10-1982 25-03-1981
JP 55133366 A	17-10-1980	JP 1351434 C JP 61016274 B	28-11-1986 28-04-1986
WO 0033837 A	15-06-2000	NONE	
WO 0017175 A	30-03-2000	AU 6251499 A	10-04-2000

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/01841

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9921555 A	06-05-1999	AU 9648098 A	17-05-1999
		EP 1027050 A	16-08-2000
		JP 11193281 A	21-07-1999
-----			